

U.S. Patent

May 15, 2007

Sheet 1 of 2

US 7,217,286 B2

FIG. 1




FIG. 1a




FIG. 2a





FIG. 2b



U.S. Patent

May 15, 2007

Sheet 2 of 2

US 7,217,286 B2

FIG. 3a





FIG. 3b



FIG. 4



US 7,217,286 B2

1

**LOCAL DELIVERY OF RAPAMYCIN FOR
TREATMENT OF PROLIFERATIVE
SEQUELAE ASSOCIATED WITH PTCA
PROCEDURES, INCLUDING DELIVERY
USING A MODIFIED STENT**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

This application is a continuation of Ser. No. 10/951,385, 10
filed Sep. 28, 2004, now pending, which in turn is a
continuation of Ser. No. 10/408,328, filed Apr. 7, 2003, now
issued as U.S. Pat. No. 6,808,536, which in turn is a
continuation of application Ser. No. 09/874,117, filed Jun. 4,
2001, now issued as U.S. Pat. No. 6,585,764, which is a 15
continuation of application Ser. No. 09/061,568, filed Apr.
16, 1998, now issued as U.S. Pat. No. 6,273,913, which in
turn claims benefit of provisional application Ser. No.
60/044,692, filed Apr. 18, 1997. The disclosures of these
prior applications are incorporated herein by reference in 20
their entirety.

FIELD OF THE INVENTION

Delivery of rapamycin locally, particularly from an intra- 25
vascular stent, directly from micropores in the stent body or
mixed or bound to a polymer coating applied on stent, to
inhibit neointimal tissue proliferation and thereby prevent
restenosis. This invention also facilitates the performance of
the stent in inhibiting restenosis.

BACKGROUND OF THE INVENTION

Re-narrowing (restenosis) of an atherosclerotic coronary 35
artery after percutaneous transluminal coronary angioplasty
(PTCA) occurs in 10–50% of patients undergoing this
procedure and subsequently requires either further angio-
plasty or coronary artery bypass graft. While the exact
hormonal and cellular processes promoting restenosis are
still being determined, our present understanding is that the 40
process of PTCA, besides opening the atherosclerotically
obstructed artery, also injures resident coronary arterial
smooth muscle cells (SMC). In response to this injury,
adhering platelets, infiltrating macrophages, leukocytes, or
the smooth muscle cells (SMC) themselves release cell 45
derived growth factors with subsequent proliferation and
migration of medial SMC through the internal elastic lamina
to the area of the vessel intima. Further proliferation and
hyperplasia of intimal SMC and, most significantly, produc-
tion of large amounts of extracellular matrix over a period of 50
3–6 months results in the filling in and narrowing of the
vascular space sufficient to significantly obstruct coronary
blood flow.

Several recent experimental approaches to preventing 55
SMC proliferation have shown promise although the
mechanisms for most agents employed are still unclear.
Heparin is the best known and characterized agent causing
inhibition of SMC proliferation both in vitro and in animal
models of balloon angioplasty-mediated injury. The mecha-
nism of SMC inhibition with heparin is still not known but 60
may be due to any or all of the following: 1) reduced
expression of the growth regulatory protooncogenes c-fos
and c-myc, 2) reduced cellular production of tissue plasmi-
nogen activator; are 3) binding and dequstration of growth
regulatory factors such as fibrovalent growth factor (FGF). 65

Other agents which have demonstrated the ability to
reduce myointimal thickening in animal models of balloon

2

vascular injury are angiopeptin (a somatostatin analog),
calcium channel blockers, angiotensin converting enzyme
inhibitors (captopril, cilazapril), cyclosporin A, trapidil (an
antianginal, antiplatelet agent), terbinafine (antifungal),
colchicine and taxol (antitubulin antiproliferatives), and
c-myc and c-myb antisense oligonucleotides. 5

Additionally, a goat antibody to the SMC mitogen platelet
derived growth factor (PDGF) has been shown to be effective
in reducing myointimal thickening in a rat model of
balloon angioplasty injury, thereby implicating PDGF
directly in the etiology of restenosis. Thus, while no therapy
has as yet proven successful clinically in preventing restenosis
after angioplasty, the in vivo experimental success of
several agents known to inhibit SMC growth suggests that
these agents as a class have the capacity to prevent clinical
restenosis and deserve careful evaluation in humans.

Coronary heart disease is the major cause of death in men
over the age of 40 and in women over the age of fifty in the
western world. Most coronary artery-related deaths are due
to atherosclerosis. Atherosclerotic lesions which limit or
obstruct coronary blood flow are the major cause of
ischemic heart disease related mortality and result in 500,
000–600,000 deaths in the United States annually. To arrest
the disease process and prevent the more advanced disease
states in which the cardiac muscle itself is compromised,
direct intervention has been employed via percutaneous
transluminal coronary angioplasty (PTCA) or coronary
artery bypass graft (CABG) PTCA is a procedure in which
a small balloon-tipped catheter is passed down a narrowed
coronary artery and then expanded to re-open the artery. It
is currently performed in approximately 250,000–300,000
patients each year. The major advantage of this therapy is
that patients in which the procedure is successful need not
undergo the more invasive surgical procedure of coronary
artery bypass graft. A major difficulty with PTCA is the
problem of post-angioplasty closure of the vessel, both
immediately after PTCA (acute reocclusion) and in the long
term (restenosis). 30

The mechanism of acute reocclusion appears to involve
several factors and may result from vascular recoil with
resulant closure of the artery and/or deposition of blood
platelets along the damaged length of the newly opened
blood vessel followed by formation of a fibrin/red blood cell
thrombus. Recently, intravascular stents have been exam- 45
ined as a means of preventing acute reclosure after PTCA.

Restenosis (chronic reclosure) after angioplasty is a more
gradual process than acute reocclusion: 30% of patients with
subtotal lesions and 50% of patients with chronic total
lesions will go on to restenosis after angioplasty. While the
exact mechanism for restenosis is still under active investi-
gation, the general aspects of the restenosis process have
been identified.

In the normal arterial wall, smooth muscle cells (SMC)
proliferate at a low rate (<0.1%/day; ref). SMC in vessel
wall exists in a *contractile* phenotype characterized by
80–90% of the cell cytoplasmic volume occupied with the
contractile apparatus. Endoplasmic reticulum, golgi bodies,
and free ribosomes are few and located in the perinuclear
region. Extracellular matrix surrounds SMC and is rich in
heparin-like glycosylaminoglycans which are believed to be
responsible for maintaining SMC in the contractile pheno-
typic state.

Upon pressure expansion of an intracoronary balloon
catheter during angioplasty, smooth muscle cells within the
arterial wall become injured. Cell derived growth factors
such as platelet derived growth factor (PDGF), basic fibro-
blast growth factor (bFGF), epidermal growth factor (EGF),

US 7,217,286 B2

3

etc. released from platelets (i.e., PDGF) adhering to the damaged arterial luminal surface, invading macrophages and/or leukocytes, or directly from SMC (i.e., BFGF) provoke a proliferation and migratory response in medial SMC. These cells undergo a phenotypic change from the contrac-

tile phenotype to a synthetic phenotype characterized by only few contractile filament bundles but extensive rough endoplasmic reticulum, golgi and free ribosomes. Proliferation/migration usually begins within 1–2 days post-injury and peaks at 2 days in the media, rapidly declining thereafter (Campbell et al., In: Vascular Smooth Muscle Cells in Culture, Campbell, J. H. and Campbell, G. R., Eds, CRC Press, Boca Ration, 1987, pp. 39–55); Clowes, A. W. and Schwartz, S. M., Circ. Res. 56:139–145, 1985).

Finally, daughter synthetic cells migrate to the intimal layer of arterial smooth muscle and continue to proliferate. Proliferation and migration continues until the damaged luminal endothelial layer regenerates at which time proliferation ceases within the intima, usually within 7–14 days postinjury. The remaining increase in intimal thickening which occurs over the next 3–6 months is due to an increase in extracellular matrix rather than cell number. Thus, SMC migration and proliferation is an acute response to vessel injury while intimal hyperplasia is a more chronic response. (Liu et al., Circulation, 79:1374–1387, 1989).

Patients with symptomatic reocclusion require either repeat PTCA or CABG. Because 30–50% of patients undergoing PTCA will experience restenosis, restenosis has clearly limited the success of PTCA as a therapeutic approach to coronary artery disease. Because SMC proliferation and migration are intimately involved with the pathophysiological response to arterial injury, prevention of SMC proliferation and migration represents a target for pharmacological intervention in the prevention of restenosis.

SUMMARY OF THE INVENTION

Novel Features and Applications to Stent Technology
Currently, attempts to improve the clinical performance of stents have involved some variation of either applying a coating to the metal, attaching a covering or membrane, or embedding material on the surface via ion bombardment. A stent designed to include reservoirs is a new approach which offers several important advantages over existing technologies.

Local Drug Delivery from a Stent to Inhibit Restenosis

In this application, it is desired to deliver a therapeutic agent to the site of arterial injury. The conventional approach has been to incorporate the therapeutic agent into a polymer material which is then coated on the stent. The ideal coating material must be able to adhere strongly to the metal stent both before and after expansion, be capable of retaining the drug at a sufficient load level to obtain the required dose, be able to release the drug in a controlled way over a period of several weeks, and be as thin as possible so as to minimize the increase in profile. In addition, the coating material should not contribute to any adverse response by the body (i.e., should be non-thrombogenic, non-inflammatory, etc.). To date, the ideal coating material has not been developed for this application.

An alternative would be to design the stent to contain reservoirs which could be loaded with the drug. A coating or membrane of biocompatible material could be applied over the reservoirs which would control the diffusion of the drug from the reservoirs to the artery wall.

One advantage of this system is that the properties of the coating can be optimized for achieving superior biocompat-

4

ibility and adhesion properties, without the addition requirement of being able to load and release the drug. The size, shape, position, and number of reservoirs can be used to control the amount of drug, and therefore the dose delivered.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will be better understood in connection with the following figures in which FIGS. 1 and 1A are top views and section views of a stent containing reservoirs as described in the present invention;

FIGS. 2a and 2b are similar views of an alternate embodiment of the stent with open ends;

FIGS. 3a and 3b are further alternate figures of a device containing a grooved reservoir; and

FIG. 4 is a layout view of a device containing a reservoir as in FIG. 3.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

Pharmacological attempts to prevent restenosis by pharmacologic means have thus far been unsuccessful and all involve systemic administration of the trial agents. Neither aspirin-dipyridamole, ticlopidine, acute heparin administration, chronic warfarin (6 months) nor methylprednisolone have been effective in preventing restenosis although platelet inhibitors have been effective in preventing acute reocclusion after angioplasty. The calcium antagonists have also been unsuccessful in preventing restenosis, although they are still under study. Other agents currently under study include thromboxane inhibitors, prostacyclin mimetics, platelet membrane receptor blockers, thrombin inhibitors and angiotensin converting enzyme inhibitors. These agents must be given systemically, however, and attainment of a therapeutically effective dose may not be possible; antiproliferative (or anti-restenosis) concentrations may exceed the known toxic concentrations of these agents so that levels sufficient to produce smooth muscle inhibition may not be reached (Lang et al., 42 Ann. Rev. Med., 127–132 (1991); Popma et al., 84 Circulation, 1426–1436 (1991)).

Additional clinical trials in which the effectiveness for preventing restenosis of dietary fish oil supplements, thromboxane receptor antagonists, cholesterol lowering agents, and serotonin antagonists has been examined have shown either conflicting or negative results so that no pharmacological agents are as yet clinically available to prevent post-angioplasty restenosis (Franklin, S. M. and Faxon, D. P., 4 Coronary Artery Disease, 2-32-242 (1993); Serruys, P. W. et al., 88 Circulation, (part 1) 1588–1601, (1993).

Conversely, stents have proven useful in preventing reducing the proliferation of restenosis. Stents, such as the stent 10 seen in layout in FIG. 4, balloon-expandable slotted metal tubes (usually but not limited to stainless steel), which when expanded within the lumen of an angioplastied coronary artery, provide structural support to the arterial wall. This support is helpful in maintaining an open path for blood flow. In two randomized clinical trials, stents were shown to increase angiographic success after PTCA, increase the stenosed blood vessel lumen and to reduce the lesion recurrence at 6 months (Serruys et al., 331 New Eng Jour. Med, 495, (1994); Fischman et al., 331 New Eng Jour. Med, 496–501 (1994). Additionally, in a preliminary trial, heparin coated stents appear to possess the same benefit of reduction in stenosis diameter at follow-up as was observed with non-heparin coated stents. Additionally, heparin coating appears to have the added benefit of producing a reduction

US 7,217,286 B2

5

in sub-acute thrombosis after stent implantation (Serruys et al., 93 *Circulation*, 412-422, (1996). Thus, 1) sustained mechanical expansion of a stenosed coronary artery has been shown to provide some measure of restenosis prevention, and 2) coating of stents with heparin has demonstrated both the feasibility and the clinical usefulness of delivering drugs to local, injured tissue off the surface of the stent.

Numerous agents are being actively studied as antiproliferative agents for use in restenosis and have shown some activity in experimental animal models. These include: heparin and heparin fragments (Clowes and Karnovsky, 265 *Nature*, 25-626, (1977); Guyton, J. R. et al. 46 *Circ. Res.*, 625-634, (1980); Clowes, A. W. and Clowes, M. M., 52 *Lab. Invest.*, 611-616, (1985); Clowes, A. W. and Clowes, M. M., 58 *Circ. Res.*, 839-845 (1986); Majesky et al., 61 *Circ. Res.*, 296-300, (1987); Snow et al., 137 *Am. J. Pathol.*, 313-330 (1990); Okada, T. et al., 25 *Neurosurgery*, 92-898, (1989) colchicine (Currier, J. W. et al., 80 *Circulation*, 11-66, (1989), taxol (ref), angiotensin converting enzyme (ACE) inhibitors (Powell, J. S. et al., 245 *Science*, 186-188 (1989), angiopeptin (Lundergan, C. F. et al., 17 *Am. J. Cardiol. (Suppl. B)*; 132B-136B (1991), Cyclosporin A (Jonasson, L. et al., 85 *Proc. Natl. Acad. Sci.*, 2303 (1988), goat-anti-rabbit PDGF antibody (Ferns, G. A. A., et al., 253 *Science*, 1129-1132 (1991), terbinafine (Nemecek, G. M. et al., 248 *J. Pharmacol. Exp. Ther.*, 1167-11747 (1989), trapidil (Liu, M. W. et al., 81 *Circulation*, 1089-1093 (1990), interferon-gamma (Hansson, G. K. and Holm, 84 *J. Circulation*, 1266-1272 (1991), steroids (Colburn, M. D. et al., 15 *J. Vasc. Surg.*, 510-518 (1992), see also Berk, B. C. et al., 17 *J. Am. Coll. Cardiol.*, 111B-117B (1991), ionizing radiation (ref), fusion toxins (ref) antisense oligonucleotides (ref), gene vectors (ref), and rapamycin (see below).

Of particular interest in rapamycin. Rapamycin is a macrolide antibiotic which blocks IL-2-mediated T-cell proliferation and possesses antiinflammatory activity. While the precise mechanism of rapamycin is still under active investigation, rapamycin has been shown to prevent the G.sub.1 to S phase progression of T-cells through the cell cycle by inhibiting specific cell cyclins and cyclin-dependent protein kinases (Siekierka, *Immunol. Res.* 13: 110-116, 1994). The antiproliferative action of rapamycin is not limited to T-cells; Marx et al. (*Circ Res* 76:412-417, 1995) have demonstrated that rapamycin prevents proliferation of both rat and human SMC in vitro while Poon et al. have shown the rat, porcine, and human SMC migratin can also be inhibited by rapamycin (*J Clin Invest* 98: 2277-2283, 1996). Thus, rapamycin is capable of inhibiting both the inflammatory response known to occur after arterial injury and stent implantation, as well as the SMC hyperproliferative response. In fact, the combined effects of rapamycin have been demonstrated to result in a diminished SMC hyperproliferative response in a rat femoral artery graft model and in both rat and porcine arterial balloon injury models (Gregory et al., *Transplantation* 55:1409-1418, 1993; Gallo et al., in press, (1997)). These observations clearly support the potential use of rapamycin in the clinical setting of post-angioplasty restenosis.

Although the ideal agent for restenosis has not yet been identified, some desired properties are clear: inhibition of local thrombosis without the risk systemic bleeding complications and continuous and prevention of the dequale of arterial injury, including local inflammation and sustained prevention smooth muscle proliferation at the site of angioplasty without serious systemic complications. Inasmuch as stents prevent at least a portion of the restenosis process, an agent which prevents inflammation and the proliferation of

6

SMC combined with a stent may provide the most efficacious treatment for post-angioplasty restenosis.

Experiments

Agents: Rapamycin (sirolimus) structural analogs (macrocyclic lactones) and inhibitors of cell-cycle progression.

Delivery Methods: These can vary:

Local delivery of such agents (rapamycin) from the struts of a stent, from a stent graft, grafts, stent cover or sheath.

Involving comixture with polymers (both degradable and nondegrading) to hold the drug to the stent or graft.

or entrapping the drug into the metal of the stent or graft body which has been modified to contain micropores or channels, as will be explained further herein.

or including covalent binding of the drug to the stent via solution chemistry techniques (such as via the Carmeda process) or dry chemistry techniques (e.g. vapour deposition methods such as rf-plasma polymerization) and combinations thereof.

Catheter delivery intravascularly from a tandem balloon or a porous balloon for intramural uptake.

Extravascular delivery by the pericardial route.

Extravascular delivery by the advential application of sustained release formulations.

Uses:

for inhibition of cell proliferation to prevent neointimal proliferation and restenosis.

prevention of tumor expansion from stents.

preventing growth of tissue into catheters and shunts inducing their failure.

1. Experimental Stent Delivery Method—Delivery from Polymer Matrix:

Solution of Rapamycin, prepared in a solvent miscible with polymer carrier solution, is mixed with solution of polymer at final concentration range 0.001 weight % to 30 weight % of drug. Polymers are biocompatible (i.e., not elicit any negative tissue reaction or promote mural thrombus formation) and degradable, such as lactone-based polyesters or copolyesters, e.g., polylactide, polycaprolactone-glycolide, polyorthoesters, polyanhydrides; poly-amino acids; polysaccharides; polyphosphazenes; poly(ether-ester) copolymers, e.g., PEO-PLLA, or blends thereof. Nonabsorbable biocompatible polymers are also suitable candidates. Polymers such as polydimethylsiloxane; poly(ethylene-vinylacetate); acrylate based polymers or copolymers, e.g., poly(hydroxyethyl methylmethacrylate, polyvinyl pyrrolidinone; fluorinated polymers such as polytetrafluoroethylene; cellulose esters.

Polymer/drug mixture is applied to the surfaces of the stent by either dip-coating, or spray coating, or brush coating or dip/spin coating or combinations thereof, and the solvent allowed to evaporate to leave a film with entrapped rapamycin.

2. Experimental Stent Delivery Method—Delivery from Microporous Depots in Stent Through a Polymer Membrane Coating:

Stent, whose body has been modified to contain micropores or channels is dipped into a solution of Rapamycin, range 0.001 wt % to saturated, in organic solvent such as acetone or methylene chloride, for sufficient time to allow solution to permeate into the pores. (The dipping solution can also be compressed to improve the loading efficiency.) After solvent has been allowed to evaporate, the stent is dipped briefly in fresh solvent to remove excess surface bound drug. A solution of polymer, chosen from any identified in the first experimental method, is applied to the

US 7,217,286 B2

7

stent as detailed above. This outer layer of polymer will act as diffusion-controller for release of drug.

3. Experimental Stent Delivery Method—Delivery Via Lysis of a Covalent Drug Tether:

Rapamycin is modified to contain a hydrolytically or enzymatically labile covalent bond for attaching to the surface of the stent which itself has been chemically derivatized to allow covalent immobilization. Covalent bonds such as ester, amides or anhydrides may be suitable for this.

4. Experimental Method—Pericardial Delivery:

A: Polymeric Sheet

Rapamycin is combined at concentration range previously highlighted, with a degradable polymer such as poly(ϵ -caprolactone-glycolid- ϵ) or non-degradable polymer, e.g., polydimethylsiloxane, and mixture cast as a thin sheet, thickness range 10.mu. to 1000.mu. The resulting sheet can be wrapped perivascularly on the target vessel. Preference would be for the absorbable polymer.

B: Conformal Coating:

Rapamycin is combined with a polymer that has a melting temperature just above 37° C., range 40°–45° C. Mixture is applied in a molten state to the external side of the target vessel. Upon cooling to body temperature the mixture solidifies conformably to the vessel wall. Both non-degradable and absorbable biocompatible polymers are suitable.

As seen in the figures it is also possible to modify currently manufactured stents in order to adequately provide the drug dosages such as rapamycin. As seen in FIGS. 1a, 2a and 3a, any stent strut 10, 20, 30 can be modified to have a certain reservoir or channel 11, 21, 31. Each of these reservoirs can be open or closed as desired. These reservoirs can hold the drug to be delivered. FIG. 4 shows a stent 40 with a reservoir 45 created at the apex of a flexible strut. Of course, this reservoir 45 is intended to be useful to deliver rapamycin or any other drug at a specific point of flexibility of the stent. Accordingly, this concept can be useful for "second generation" type stents.

In any of the foregoing devices, however, it is useful to have the drug dosage applied with enough specificity and

8

enough concentration to provide an effective dosage in the lesion area. In this regard, the reservoir size in the stent struts must be kept at a size of about 0.0005" to about 0.003". Then, it should be possible to adequately apply the drug dosage at the desired location and in the desired amount.

These and other concepts will be disclosed herein. It would be apparent to the reader that modifications are possible to the stent or the drug dosage applied. In any event, however, the any obvious modifications should be perceived to fall within the scope of the invention which is to be realized from the attached claims and their equivalents.

What is claimed:

1. A device comprising a metallic stent, a biocompatible, nonabsorbable polymeric carrier, and a therapeutic agent, wherein:

said polymeric carrier comprises an acrylate-based polymer or copolymer, a fluorinated polymer, or a mixture thereof, and

said therapeutic agent is rapamycin, or a macrocyclic lactone analog thereof, and is present in an amount effective to inhibit neointimal proliferation.

2. The device according to claim 1 wherein said therapeutic agent is a macrocyclic lactone analog of rapamycin.

3. The device according to claim 1 that provides a controlled release of said therapeutic agent over a period of several weeks.

4. The device according to claim 2 that provides a controlled release of said therapeutic agent over a period of several weeks.

5. A method of inhibiting neointimal proliferation in a coronary artery resulting from percutaneous transluminal coronary angioplasty comprising implanting a device according to any one of claims 1 to 4 in the lumen of said coronary artery.

* * * * *



US007223286B2

(12) **United States Patent**
Wright et al.

(10) **Patent No.:** **US 7,223,286 B2**(45) **Date of Patent:** ***May 29, 2007**

(54) **LOCAL DELIVERY OF RAPAMYCIN FOR TREATMENT OF PROLIFERATIVE SEQUELAE ASSOCIATED WITH PTCA PROCEDURES, INCLUDING DELIVERY USING A MODIFIED STENT**

(58) **Field of Classification Search** 623/1.42–1.48;
427/2.1–2.31
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

861,659 A	7/1907	Johnston	464/147
3,051,677 A	8/1962	Rexford	522/156
3,279,996 A	10/1966	Long et al.	424/424
3,526,005 A	9/1970	Bokros	623/11.11
3,599,641 A	8/1971	Sheridan	604/256
3,657,744 A	4/1972	Ersek	128/898

(75) **Inventors:** **Carol Wright**, Somerset, NJ (US);
Gerard H. Llanos, Stewartsville, NJ (US); **Ronald Rakos**, Neshanic Station, NJ (US); **Kristin King**, Mahwah, NJ (US); **Robert Falotico**, Bell Mead, NJ (US)

(73) **Assignee:** **Cordis Corporation**, Miami Lakes, FL (US)

(*) **Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 265 days.

This patent is subject to a terminal disclaimer.

(Continued)

FOREIGN PATENT DOCUMENTS

DE 3205942 A1 9/1983

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 07/819,314, filed Jan. 9, 1992, Morris.

(Continued)

Primary Examiner—Suzette Gherbi

(74) *Attorney, Agent, or Firm*—Woodcock Washburn LLP

(57) **ABSTRACT**

Methods of preparing intravascular stents with a polymeric coating containing macrocyclic lactone (such as rapamycin or its analogs), stents and stent graphs with such coatings, and methods of treating a coronary artery with such devices. The macrocyclic lactone-based polymeric coating facilitates the performance of such devices in inhibiting restenosis.

(21) **Appl. No.:** **10/951,385**

(22) **Filed:** **Sep. 28, 2004**

(65) **Prior Publication Data**

US 2005/0085902 A1 Apr. 21, 2005

Related U.S. Application Data


(63) Continuation of application No. 10/408,328, filed on Apr. 7, 2003, now Pat. No. 6,808,536, which is a continuation of application No. 09/874,117, filed on Jun. 4, 2001, now Pat. No. 6,585,764, which is a continuation of application No. 09/061,568, filed on Apr. 16, 1998, now Pat. No. 6,273,913.

(60) Provisional application No. 60/044,692, filed on Apr. 18, 1997.

(51) **Int. Cl.**
A61F 2/06 (2006.01)

(52) **U.S. Cl.** 623/1.42

77 Claims, 2 Drawing Sheets



US 7,223,286 B2

Page 2

U.S. PATENT DOCUMENTS

3,744,596 A	7/1973	Sander	188/203	5,049,132 A	9/1991	Shaffer et al.	604/101.02
3,779,805 A	12/1973	Alsberg	427/105	5,049,403 A	9/1991	Larm et al.	427/2.1
3,929,992 A	12/1975	Sehgal et al.	424/122	5,053,048 A	10/1991	Pinchuk	623/1.43
3,932,627 A	1/1976	Margraf	514/56	5,059,166 A	10/1991	Fischell et al.	600/3
3,948,254 A	4/1976	Zaffaroni	128/833	5,061,275 A	10/1991	Wallsten et al.	623/1.22
3,952,334 A	4/1976	Bokros et al.	623/11.11	5,061,750 A	10/1991	Feijen et al.	525/54.1
3,968,800 A	7/1976	Vilasi	606/198	5,064,435 A	11/1991	Porter	623/23.7
4,069,307 A	1/1978	Higuchi et al.	424/432	5,092,877 A	3/1992	Pinchuk	128/898
4,076,285 A	2/1978	Martinez	285/332	5,102,417 A	4/1992	Palmaz	606/195
4,292,965 A	10/1981	Nash et al.	128/833	5,104,404 A	4/1992	Wolff	623/1.16
4,299,226 A	11/1981	Banka	604/509	5,116,365 A	5/1992	Hillstead	623/1.15
4,300,244 A	11/1981	Bokros	623/1.13	5,122,154 A	6/1992	Rhodes	623/1.13
4,312,920 A	1/1982	Pierce et al.	428/425.5	5,131,908 A	7/1992	Dardik et al.	600/36
4,321,711 A	3/1982	Mano	623/1.43	5,133,732 A	7/1992	Wiktor	623/1.22
4,323,071 A	4/1982	Simpson et al.	606/194	5,134,192 A	7/1992	Feijen et al.	525/54.1
4,390,599 A	6/1983	Broyles	428/597	5,135,536 A	8/1992	Hillstead	606/195
4,413,359 A	11/1983	Akiyama et al.	623/23.72	5,163,952 A	11/1992	Froix	623/1.18
4,423,183 A	12/1983	Close	524/546	5,163,958 A	11/1992	Pinchuk	623/23.49
4,441,216 A	4/1984	Ionescu et al.	623/2.19	5,171,217 A	12/1992	March et al.	604/507
4,503,569 A	3/1985	Dotter	623/1.19	5,171,262 A	12/1992	MacGregor	623/1.15
4,512,338 A	4/1985	Balko et al.	606/108	5,176,660 A	1/1993	Truckai	604/527
4,550,447 A	11/1985	Seiler, Jr. et al.	623/1.32	5,176,972 A	1/1993	Bloom et al.	430/14
4,553,545 A	11/1985	Maass et al.	606/198	5,178,618 A	1/1993	Kandarpa	606/28
4,560,374 A	12/1985	Hammerslag	604/509	5,180,366 A	1/1993	Woods	604/96.01
4,562,596 A	1/1986	Kornberg	623/1.32	5,182,317 A	1/1993	Winters et al.	523/112
4,565,740 A	1/1986	Golander et al.	428/409	5,185,408 A	2/1993	Tang et al.	525/415
4,580,568 A	4/1986	Gianturco	606/198	5,192,307 A	3/1993	Wall	623/1.2
4,613,665 A	9/1986	Larm	536/20	5,195,984 A	3/1993	Schatz	623/1.2
4,642,111 A	2/1987	Sakamoto et al.	424/492	5,213,576 A	5/1993	Abiuso et al.	604/103.01
4,655,771 A	4/1987	Wallsten	623/1.22	5,213,898 A	5/1993	Larm et al.	428/422
4,656,083 A	4/1987	Hoffman et al.	442/123	5,217,483 A	6/1993	Tower	623/1.15
4,676,241 A	6/1987	Webb et al.	128/207.14	5,222,971 A	6/1993	Willard et al.	606/198
4,678,466 A	7/1987	Rosenwald	424/427	5,226,913 A	7/1993	Pinchuk	140/71 R
4,687,482 A	8/1987	Hanson	623/1.49	5,234,456 A	8/1993	Silvestrini	623/1.2
4,689,046 A	8/1987	Bokros	623/2.31	5,246,445 A	9/1993	Yachia et al.	623/1.2
4,731,054 A	3/1988	Billeter et al.	604/93.01	5,258,020 A	11/1993	Froix	128/898
4,733,665 A	3/1988	Palmaz	606/108	5,258,021 A	11/1993	Duran	623/2.3
4,739,762 A	4/1988	Palmaz	623/1.11	5,262,451 A	11/1993	Winters et al.	523/112
4,740,207 A	4/1988	Kreamer	623/1.15	5,266,073 A	11/1993	Wall	623/1.2
4,749,585 A	6/1988	Greco et al.	428/422	5,272,012 A	12/1993	Opolski	428/423.1
4,753,652 A	6/1988	Langer et al.	623/1.42	4,733,665 A	1/1994	Palmaz	606/108
4,760,849 A	8/1988	Kropf	606/191	5,275,622 A	1/1994	Lazarus et al.	623/1.11
4,768,507 A	9/1988	Fischell et al.	623/1.11	5,282,823 A	2/1994	Schwartz et al.	623/1.22
4,776,337 A	10/1988	Palmaz	623/1.11	5,282,824 A	2/1994	Gianturco	623/1.13
4,786,500 A	11/1988	Wong	424/422	5,283,257 A	2/1994	Gregory et al.	514/458
4,787,899 A	11/1988	Lazarus	623/1.11	5,288,711 A	2/1994	Mitchell et al.	424/122
4,800,882 A	1/1989	Gianturco	606/194	5,290,305 A	3/1994	Inoue	623/1.2
4,810,784 A	3/1989	Larm	536/20	5,292,331 A	3/1994	Boneau	623/1.16
4,856,516 A	8/1989	Hillstead	606/194	5,292,802 A	3/1994	Rhee et al.	525/54.1
4,871,357 A	10/1989	Hsu et al.	604/266	5,304,121 A	4/1994	Sahatjian	604/509
4,872,867 A	10/1989	Joh	604/269	5,304,200 A	4/1994	Spaulding	623/1.16
4,876,109 A	10/1989	Mayer et al.	604/269	5,306,250 A	4/1994	March et al.	604/104
4,886,062 A	12/1989	Wiktor	606/194	5,308,862 A	5/1994	Ohlstein	514/411
4,907,336 A	3/1990	Gianturco	29/515	5,308,889 A	5/1994	Rhee et al.	523/113
4,916,193 A	4/1990	Tang et al.	525/413	5,314,444 A	5/1994	Gianturco	606/195
4,954,126 A	9/1990	Wallsten	600/36	5,314,472 A	5/1994	Fontaine	623/1.22
4,969,458 A	11/1990	Wiktor	623/1.11	5,328,471 A	7/1994	Slepian	604/101.03
4,990,131 A	2/1991	Dardik et al.	600/36	5,334,301 A	8/1994	Heinke et al.	204/267
4,990,155 A	2/1991	Wilko	606/191	5,336,518 A	8/1994	Narayanan et al.	427/470
4,994,071 A	2/1991	MacGregor	606/194	5,338,770 A	8/1994	Winters et al.	523/112
4,994,298 A	2/1991	Yasuda	427/490	5,342,348 A	8/1994	Kaplan	604/891.1
4,998,923 A	3/1991	Samson et al.	606/194	5,342,387 A	8/1994	Summers	606/198
5,015,253 A	5/1991	MacGregor	623/1.15	5,342,621 A	8/1994	Eury	606/198
5,019,090 A	5/1991	Pinchuk	623/1.15	5,354,257 A	10/1994	Roubin et al.	600/7
5,019,096 A	5/1991	Fox, Jr. et al.	600/36	5,354,308 A	10/1994	Simon et al.	623/1.15
5,029,877 A	7/1991	Fedeli	277/354	5,356,433 A	10/1994	Rowland et al.	424/422
5,034,265 A	7/1991	Hoffman et al.	442/126	5,366,504 A	11/1994	Andersen et al.	623/1.5
5,035,706 A	7/1991	Giantureo et al.	606/198	5,368,566 A	11/1994	Crocker	604/101.02
5,041,100 A	8/1991	Rowland et al.	604/265	5,370,683 A	12/1994	Fontaine	623/1.22
5,041,126 A	8/1991	Gianturco	623/1.15	5,370,691 A	12/1994	Samson	623/1.22
5,047,020 A	9/1991	Hsu	604/266	5,375,612 A	12/1994	Cottenceau et al.	128/899
				5,376,112 A	12/1994	Duran	623/1.26
				5,378,475 A	1/1995	Smith et al.	424/473

US 7,223,286 B2

Page 3

5,380,299 A	1/1995	Fearnot et al.	604/265	5,605,696 A	2/1997	Eury et al.	424/423
5,382,261 A	1/1995	Palmaz	606/158	5,607,463 A	3/1997	Schwartz et al.	623/1.44
5,383,853 A	1/1995	Jung et al.	604/103.04	5,607,475 A	3/1997	Cahalan et al.	424/423
5,383,928 A	1/1995	Scott et al.	623/1.12	5,609,629 A	3/1997	Fearnot et al.	623/1.42
5,387,235 A	2/1995	Chuter	623/1.11	5,616,608 A	4/1997	Kinsella et al.	514/449
5,389,106 A	2/1995	Tower	623/1.15	5,620,984 A	4/1997	Bianco et al.	514/263.36
5,393,772 A	2/1995	Yue et al.	514/410	5,621,102 A	4/1997	Bianco et al.	544/267
5,395,390 A	3/1995	Simon et al.	623/1.18	5,622,975 A	4/1997	Singh et al.	514/324
5,397,355 A	3/1995	Marin et al.	623/1.2	5,624,411 A	4/1997	Tuch	604/265
5,399,352 A	3/1995	Hanson	424/423	5,628,785 A	5/1997	Schwartz et al.	128/898
5,403,341 A	4/1995	Solar	606/198	5,629,077 A	5/1997	Turnlund et al.	623/1.15
5,405,377 A	4/1995	Cragg	623/1.2	5,629,315 A	5/1997	Bianco et al.	514/263.36
5,409,696 A	4/1995	Narayanan et al.	424/78.17	5,632,763 A	5/1997	Glastra	623/1.15
5,411,549 A	5/1995	Peters	623/1.15	5,632,771 A	5/1997	Boatman et al.	623/1.15
5,415,619 A	5/1995	Lee et al.	600/36	5,632,776 A	5/1997	Kurumatani et al.	424/423
5,417,969 A	5/1995	Hsu et al.	424/78.27	5,632,840 A	5/1997	Campbell	156/196
5,419,760 A	5/1995	Narciso, Jr.	604/8	5,635,201 A	6/1997	Fabo	424/443
D359,802 S	6/1995	Fontaine	D24/155	5,637,113 A	6/1997	Tartaglia et al.	623/1.42
5,421,955 A	6/1995	Lau et al.	216/48	5,643,312 A	7/1997	Fischell et al.	623/1.15
5,423,885 A	6/1995	Williams	623/1.17	5,643,939 A	7/1997	Ohlstein	514/411
5,429,618 A	7/1995	Keogh	604/266	5,646,160 A	7/1997	Morris et al.	514/291
5,429,634 A	7/1995	Narciso, Jr.	604/890.1	5,648,357 A	7/1997	Bianco et al.	514/263.36
5,439,446 A	8/1995	Barry	604/103.01	5,649,952 A	7/1997	Lam	623/1.15
5,441,515 A	8/1995	Khosravi et al.	606/194	5,649,977 A	7/1997	Campbell	623/1.15
5,441,516 A	8/1995	Wang et al.	606/198	5,651,174 A	7/1997	Schwartz et al.	29/527.2
5,441,947 A	8/1995	Dodge et al.	514/179	5,652,243 A	7/1997	Bianco et al.	514/263.36
5,443,458 A	8/1995	Evry	604/891.1	5,653,747 A	8/1997	Dereume	623/1.54
5,443,477 A	8/1995	Marin et al.	606/198	5,653,992 A	8/1997	Bezawada et al.	424/426
5,443,496 A	8/1995	Schwartz et al.	623/1.16	5,662,609 A	9/1997	Slepian	604/101.03
5,443,498 A	8/1995	Fontaine	623/1.17	5,665,591 A *	9/1997	Sonenshein et al.	435/375
5,443,500 A	8/1995	Sigwart	623/1.17	5,665,728 A	9/1997	Morris et al.	424/122
5,447,724 A	9/1995	Helmus et al.	424/426	5,667,764 A	9/1997	Kopia et al.	424/1.45
5,449,372 A	9/1995	Schmaltz et al.	606/198	5,669,924 A	9/1997	Shaknovich	623/1.11
5,449,373 A	9/1995	Pinchasik et al.	606/198	5,670,506 A	9/1997	Leigh et al.	514/141
5,449,382 A	9/1995	Dayton	623/1.15	5,672,638 A	9/1997	Verhoeven et al.	523/112
5,464,450 A	11/1995	Buscemi et al.	632/1.2	5,674,242 A	10/1997	Phan et al.	606/198
5,464,540 A	11/1995	Friesen et al.	210/640	5,679,400 A	10/1997	Tuch	427/2.14
5,464,650 A	11/1995	Berg et al.	427/2.3	5,679,659 A	10/1997	Verhoeven et al.	514/56
5,474,563 A	12/1995	Myler et al.	606/108	5,684,061 A	11/1997	Ohnishi et al.	523/114
5,486,357 A	1/1996	Narayanan	424/78.17	5,691,311 A	11/1997	Maraganore et al.	514/12
5,496,365 A	3/1996	Sgro	623/1.2	5,693,085 A	12/1997	Buirge et al.	623/1.13
5,500,013 A	3/1996	Buscemi et al.	623/1.22	5,697,967 A	12/1997	Dinh et al.	128/898
5,510,077 A	4/1996	Dinh et al.	264/485	5,697,971 A	12/1997	Fischell et al.	623/1.15
5,512,055 A	4/1996	Domb et al.	604/265	5,700,286 A	12/1997	Tartaglia et al.	623/1.15
5,516,781 A	5/1996	Morris et al.	514/291	5,707,385 A	1/1998	Williams	606/192
5,519,042 A	5/1996	Morris et al.	514/378	5,709,874 A	1/1998	Hanson et al.	424/423
5,523,092 A	6/1996	Hanson et al.	424/423	5,713,949 A	2/1998	Jayaraman	623/1.12
5,527,354 A	6/1996	Fontaine et al.	623/1.17	5,716,981 A	2/1998	Hunter et al.	514/449
5,545,208 A	8/1996	Wolff et al.	623/1.22	5,725,549 A	3/1998	Lam	623/1.15
5,551,954 A	9/1996	Buscemi et al.	623/1.15	5,725,567 A	3/1998	Wolff et al.	623/1.42
5,554,182 A	9/1996	Dinh et al.	600/36	5,728,150 A	3/1998	McDonald et al.	623/1.15
5,554,954 A	9/1996	Takahashi	327/546	5,728,420 A	3/1998	Keogh	427/2.12
5,556,413 A	9/1996	Lam	623/1.2	5,731,326 A	3/1998	Hart et al.	514/323
5,562,922 A	10/1996	Lambert	424/486	5,733,327 A	3/1998	Igaki et al.	623/1.5
5,563,146 A	10/1996	Morris	514/291	5,733,920 A	3/1998	Mansuri et al.	514/337
5,569,197 A	10/1996	Helmus	604/102.02	5,733,925 A	3/1998	Kunz et al.	514/449
5,569,295 A	10/1996	Lam	606/198	5,735,897 A	4/1998	Buirge	623/1.15
5,569,462 A	10/1996	Martinson et al.	424/423	5,739,138 A	4/1998	Bianco et al.	514/263.36
5,569,463 A	10/1996	Helmus et al.	424/426	5,755,734 A	5/1998	Richter et al.	606/194
5,571,089 A	11/1996	Crocker		5,755,772 A	5/1998	Evans et al.	128/898
5,571,166 A	11/1996	Dinh et al.	128/898	5,759,205 A	6/1998	Valentini	433/173
5,574,059 A	11/1996	Regunathan et al.	514/397	5,769,883 A	6/1998	Buscemi et al.	623/1.42
5,575,818 A	11/1996	Pinchuk	623/1.15	5,776,184 A	7/1998	Tuch	623/1.11
5,578,075 A	11/1996	Dayton	623/1.15	5,780,476 A	7/1998	Underiner et al.	514/263.36
5,580,873 A	12/1996	Bianco et al.	514/263.36	5,782,908 A	7/1998	Cahalan et al.	623/1.13
5,580,874 A	12/1996	Bianco et al.	514/263.36	5,788,979 A	8/1998	Alt et al.	424/426
5,591,140 A	1/1997	Narayanan et al.	604/269	5,792,106 A	8/1998	Mische	604/103.01
5,591,197 A	1/1997	Orth et al.		5,792,772 A	8/1998	Bianco et al.	514/263.36
5,591,224 A	1/1997	Schwartz et al.	623/1.22	5,798,372 A	8/1998	Davies et al.	514/356
5,591,227 A	1/1997	Dinh et al.	623/1.22	5,799,384 A	9/1998	Schwartz et al.	29/458
5,599,352 A	2/1997	Dinh et al.	128/898	5,800,507 A	9/1998	Schwartz	623/1.11
5,603,722 A	2/1997	Phan et al.	623/1.18	5,800,508 A	9/1998	Goicoechea et al.	623/1.15
5,604,283 A	2/1997	Wada et al.	524/236	5,807,861 A	9/1998	Klein et al.	514/263.35

US 7,223,286 B2

Page 4

5,811,447 A	9/1998	Kunz et al.	514/411	6,258,121 B1	7/2001	Yang et al.	623/1.46
5,820,917 A	10/1998	Tuch 427/2.1		6,268,390 B1	7/2001	Kunz 514/411	
5,820,918 A	10/1998	Ronan et al.	427/2.1	6,273,913 B1	8/2001	Wright et al.	623/1.42
5,824,048 A	10/1998	Tuch 128/898		6,284,305 B1	9/2001	Ding et al.	427/2.28
5,824,049 A	10/1998	Ragheb et al.	623/1.44	6,287,320 B1	9/2001	Slepian 606/194	
5,827,587 A	10/1998	Fukushi 428/36.6		6,287,628 B1	9/2001	Hossainy et al.	427/2.3
5,833,651 A	11/1998	Donovan et al.	604/509	6,299,604 B1	10/2001	Ragheb et al.	604/265
5,837,008 A	11/1998	Berg et al. 427/2.21		6,306,144 B1	10/2001	Sydney et al.	606/108
5,837,313 A	11/1998	Ding et al. 427/2.21		6,306,166 B1	10/2001	Barry et al.	623/1.46
5,843,120 A	12/1998	Israel et al.		6,306,176 B1	10/2001	Whitbourne 623/23.59	
5,843,166 A	12/1998	Lentz et al.		6,306,421 B1	10/2001	Kunz et al. 424/423	
5,843,172 A	12/1998	Yan 623/1.42		6,309,380 B1	10/2001	Larson et al.	604/502
5,849,034 A	12/1998	Schwartz 606/36		6,309,660 B1	10/2001	Hsu et al. 424/425	
5,851,217 A	12/1998	Wolff et al. 606/191		6,313,264 B1	11/2001	Caggiano et al.	530/350
5,851,231 A	12/1998	Wolff et al. 623/1.42		6,316,018 B1	11/2001	Ding et al. 424/423	
5,858,990 A	1/1999	Walsh 514/44		6,335,029 B1	1/2002	Kamath et al.	424/423
5,861,027 A	1/1999	Trapp 623/1.15		6,358,556 B1	3/2002	Ding et al. 427/2.24	
5,865,814 A	2/1999	Tuch 623/1.15		6,369,039 B1	4/2002	Palasis et al.	424/93.2
5,871,535 A	2/1999	Wolff et al. 128/898		6,379,382 B1	4/2002	Yang 623/1.42	
5,873,904 A	2/1999	Ragheb et al. 623/1.13		6,387,121 B1	5/2002	Alt 623/1.15	
5,876,433 A	3/1999	Lunn 623/1.15		6,403,635 B1	6/2002	Kinsella et al.	514/449
5,877,224 A	3/1999	Brocchini et al.	514/772.2	6,407,067 B1	6/2002	Schafer 514/19	
5,879,697 A	3/1999	Ding et al. 424/422		6,517,858 B1	2/2003	Haberbosch et al.	424/424
5,882,335 A	3/1999	Leone et al. 604/103.02		6,517,889 B1	2/2003	Jayaraman 427/2.24	
5,891,108 A	4/1999	Leone et al.		6,545,097 B2	4/2003	Pinchuk et al.	525/240
5,893,840 A	4/1999	Hull et al. 604/103.02		6,585,764 B2	7/2003	Wright et al.	623/1.42
5,897,911 A	4/1999	Loeffler 427/2.25		6,620,194 B2	9/2003	Ding et al. 623/1.43	
5,900,246 A	5/1999	Lambert 424/429		6,746,773 B2	6/2004	Llanos et al. 428/421	
5,902,266 A	5/1999	Leone et al. 604/509		6,776,796 B2	8/2004	Llanos et al. 623/1.46	
5,916,910 A	6/1999	Lai 514/423		6,808,536 B2	10/2004	Wright et al.	623/1.42
5,922,393 A	7/1999	Jayaraman 427/2.3		2001/0007083 A1	7/2001	Roorda 623/1.15	
5,932,243 A	8/1999	Fricker et al.		2001/0029351 A1	10/2001	Falotico et al.	604/103.02
5,932,299 A	8/1999	Katoot 427/508		2001/0029660 A1	10/2001	Johnson 29/557	
5,932,580 A	8/1999	Levitzi et al. 181/152		2001/0032014 A1	10/2001	Yang et al. 623/1.15	
5,951,586 A	9/1999	Berg et al. 606/198		2001/0034363 A1	10/2001	Li et al. 514/449	
5,957,971 A	9/1999	Schwartz 623/1.15		2001/0037145 A1	11/2001	Guruwaiya et al.	623/1.15
5,968,091 A	10/1999	Pinchuk et al. 623/1.16		2002/0010418 A1	1/2002	Lary et al. 604/101.04	
5,972,027 A	10/1999	Johnson 424/145.1		2002/0032477 A1	3/2002	Helmus et al.	623/1.2
5,976,534 A	11/1999	Hart et al. 424/145.1		2002/0041899 A1	4/2002	Chudzik et al.	424/487
5,977,163 A	11/1999	Li et al. 514/449		2002/0061326 A1	5/2002	Li et al. 424/424	
5,980,553 A	11/1999	Gray et al. 623/1.15		2002/0068969 A1	6/2002	Shanley et al.	623/1.16
5,980,566 A	11/1999	Alt et al. 623/23.7		2002/0071902 A1	6/2002	Ding et al. 427/2.24	
5,980,972 A	11/1999	Ding 427/2.24		2002/0082680 A1	6/2002	Shanley et al.	623/1.16
5,981,568 A	11/1999	Kunz et al. 514/411		2002/0082685 A1	6/2002	Sirhan et al. 623/1.42	
5,985,307 A	11/1999	Hanson et al. 424/423		2002/0091433 A1	7/2002	Ding et al. 623/1.2	
5,997,468 A	12/1999	Wolff et al. 606/36		2002/0095114 A1	7/2002	Palasis 604/96.01	
6,004,346 A	12/1999	Wolff et al. 623/23.71		2002/0099438 A1	7/2002	Furst 623/1.16	
6,015,432 A	1/2000	Rakos et al. 623/1.13		2002/0103526 A1	8/2002	Steinke 623/1.11	
6,039,721 A	3/2000	Johnson et al. 604/508		2002/0119178 A1	8/2002	Levesque et al.	424/423
6,059,813 A	5/2000	Vrba et al. 606/198		2002/0123505 A1	9/2002	Molliston et al.	514/291
6,071,305 A	6/2000	Brown et al. 623/1.43		2002/0127327 A1	9/2002	Schwartz et al.	427/2.15
6,074,659 A	6/2000	Kunz et al. 424/423		2002/0133222 A1	9/2002	Das 623/1.16	
6,080,190 A	6/2000	Schwartz 623/1.22		2002/0133224 A1	9/2002	Bajgar et al.	623/1.39
6,096,070 A	8/2000	Ragheb et al. 623/1.39		2002/0165608 A1	11/2002	Llanos 604/500	
6,120,536 A	9/2000	Ding et al. 623/1.43		2002/0193475 A1	12/2002	Hossainy et al.	524/113
6,120,847 A	9/2000	Yang et al.		2003/0065377 A1	4/2003	Davila et al.	604/265
6,136,798 A	10/2000	Cody et al. 514/141		2003/0216699 A1	11/2003	Falotico 604/265	
6,140,127 A	10/2000	Sprague 435/395		2004/0049265 A1	3/2004	Ding et al. 623/1.42	
6,146,358 A	11/2000	Rowe 604/103		2004/0243097 A1	12/2004	Falotico et al.	604/500
6,153,252 A	11/2000	Hossainy et al. 427/2.3		2004/0260268 A1	12/2004	Falotico et al.	604/500
6,159,488 A	12/2000	Nagler et al. 424/423		2005/0002986 A1	1/2005	Falotico et al.	424/426
6,171,232 B1	1/2001	Papandreou et al.	600/36	2005/0004663 A1	1/2005	Llanos et al.	623/1.46
6,171,609 B1	1/2001	Kunz 424/422		2005/0033261 A1	2/2005	Falotico et al.	604/500
6,177,272 B1	1/2001	Nabel et al. 435/320.1		2005/0106210 A1	5/2005	Ding et al. 424/423	
6,179,817 B1	1/2001	Zhong 604/265		2005/0187611 A1	8/2005	Ding et al. 623/1.15	
6,193,746 B1	2/2001	Strecker 623/1.13		2005/0208200 A1	9/2005	Ding et al. 427/2.25	
6,214,901 B1	4/2001	Chudzik et al. 523/113		2006/0088654 A1	4/2006	Ding et al. 427/2.21	
6,225,346 B1	5/2001	Tang et al. 514/523		2006/0089705 A1	4/2006	Ding et al. 623/1.15	
6,240,616 B1	6/2001	Yan 29/527.2					
6,245,537 B1	6/2001	Williams et al.	435/135				
6,251,920 B1	6/2001	Grainger et al.	514/319				
6,254,632 B1	7/2001	Wu et al. 623/1.15					
6,254,634 B1	7/2001	Anderson et al.	623/1.42				

FOREIGN PATENT DOCUMENTS

DE	19723723 A1	12/1998
EP	0 145 166 A2	6/1985

US 7,223,286 B2

Page 5

EP 0 177 330 A2 4/1986
 EP 0 183 372 A1 6/1986
 EP 0 221 570 A2 5/1987
 EP 0 421 729 A2 4/1991
 EP 0 540 290 A2 5/1993
 EP 0 568 310 A1 11/1993
 EP 0 604 022 A1 6/1994
 EP 0 621 015 A1 10/1994
 EP 0 623 354 A1 11/1994
 EP 0 734 698 A2 3/1996
 EP 0 712 615 A1 5/1996
 EP 0 716 836 A1 6/1996
 EP 0 734 721 A2 10/1996
 EP 0 747 069 A2 12/1996
 EP 0 761 251 A1 3/1997
 EP 0 800 801 A1 10/1997
 EP 0 540 290 B1 1/1998
 EP 0 830 853 A1 3/1998
 EP 0 815 803 A1 7/1998
 EP 0 850 651 A2 7/1998
 EP 0 938 878 A2 9/1999
 EP 0 938 878 A3 9/1999
 EP 0 950 386 A2 10/1999
 EP 0 968 688 A1 1/2000
 EP 0 633 032 B1 2/2001
 EP 1 192 957 A2 4/2002
 EP 1 588 726 A1 10/2005
 EP 1 588 727 A1 10/2005
 FR 566 807 A1 4/1992
 GB 0 662 307 A2 12/1951
 GB 1 205 743 A 9/1970
 GB 2 135 585 A 9/1984
 SU 660689 5/1979
 SU 1457921 2/1989
 WO 89/03232 A1 4/1989
 WO 91/12779 A1 9/1991
 WO 92/15286 A1 9/1992
 WO 94/01056 A1 1/1994
 WO 94/21308 A1 9/1994
 WO 94/21309 A1 9/1994
 WO 94/24961 A1 11/1994
 WO 96/00272 A1 1/1996
 WO 96/26689 A1 9/1996
 WO 96/32907 A1 10/1996
 WO 96/34580 A1 11/1996
 WO 97/25000 A1 7/1997
 WO 97/33534 A1 9/1997
 WO 98/08463 A1 3/1998
 WO 98/13344 A1 4/1998
 WO 98/19628 A1 5/1998
 WO 98/23228 A1 6/1998
 WO 98/23244 A1 6/1998
 WO 98/34669 A1 8/1998
 WO 98/36784 A1 8/1998
 WO 98/47447 A1 10/1998
 WO 98/56312 A1 12/1998
 WO 00/21584 A1 4/2000
 WO 00/27445 A1 5/2000
 WO 00/27455 A1 5/2000
 WO 00/32255 A1 6/2000
 WO 00/38754 A1 7/2000
 WO 01/87342 A2 11/2001
 WO 01/87372 A1 11/2001
 WO 01/87373 A1 11/2001
 WO 01/87376 A1 11/2001
 WO 02/26139 A1 4/2002
 WO 02/26271 A1 4/2002
 WO 02/26280 A1 4/2002
 WO 02/26281 A1 4/2002
 WO 03/015664 A1 2/2003

WO 03/057218 A1 7/2003

OTHER PUBLICATIONS

U.S. Appl. No. 08/424,884, filed Apr. 19, 1995, Helmus et al.
 U.S. Appl. No. 08/526,273, filed Sep. 11, 1995, Ding.
 U.S. Appl. No. 08/730,542, filed Oct. 11, 1996, Helmus.
 U.S. Appl. No. 09/575,480, filed May 19, 2000, Kopia.
 U.S. Appl. No. 10/431,059, filed May 7, 2003, Falotico
 Abraham, R. T., "Mammalian target of rapamycin: Immunosuppressive drugs offer new insight into cell growth regulation," *Progress in Inflammation Research*, 2000, Switzerland.
 Alvarado, R. et al., "Evaluation of Polymer-coated Balloon-expandable Stents in Bile Ducts," *Radiology*, 1989, 170, 975-978.
 Bailey et al., "Polymer Coating of Palmaz-Schatz Stent Attenuates Vascular Spasm after Stent Placement," *Circulation*, 82:III-541 (1990).
 Berk, B. C. et al., "Pharmacologic Roles of Heparin and Glucocorticoids to Prevent Restenosis After Coronary Angioplasty," *JACC*, May 1991, 17(6), 111B-117B.
 Bertram, P. G. et al., "The 14-3-3 proteins positively regulate rapamycin-sensitive signaling," *Current Biology*, 1998, 8, 1259-1267.
 Biomaterials Science (B.D. Ratner, Ed.), Academic Press, New York, NY, pp. 228-238, 1996.
 Campbell, G. R. et al., "Phenotypic Modulation of Smooth Muscle Cells in Primary Culture, Vascular Smooth Muscle Cells in Culture," *CRC Press*, 1987, 39-55.
 Chang, M. W. et al., "Adenovirus-mediated Over-expression of the Cyclin/Cyclin-dependent Kinase inhibitor, p21 inhibits Vascular Smooth Muscle Cell Proliferation and Neointima Formation in the Rat Carotid Artery Model of Balloon Angioplasty," *J. Clin. Invest.*, 1995, 96, 2260-2268.
 Chung, J. et al., "Rapamycin-FKBP specifically blocks growth-dependent activation of and signaling by the 70 kd S6 protein kinases," *Cell*, Jun. 26, 1992, 69(7), 1227-1236.
 Clowes, A. W. et al., "Kinetics of cellular proliferation after arterial injury. IV. Heparin inhibits rat smooth muscle mitogenesis and migration," *Circ. Res.*, 1986, 58(6), 839-845.
 Clowes, A. W. et al., Kinetics of Cellular Proliferation after Arterial Injury, *Laboratory Investigation*, 1985, 52(6), 611-616.
 Clowes, A. W. et al., "Significance of quiescent smooth muscle migration in the injured rat carotid artery," *Circ. Res.* 1985, 56(1), 139-145.
 Clowes, A. W., "Suppression by heparin of smooth muscle cell proliferation in injured arteries," *Nature*, 1977, 265(5595), 625-626.
 Colburn, M. D. et al., "Dose responsive suppression of myointimal hyperplasia by dexamethasone," *J. Vasc. Surg.*, 1992, 15, 510-518.
 Curier, J. W. et al., "Colchicine Inhibits Restenosis After Iliac Angioplasty in the Artherosclerotic Rabbit," *Circ.*, 1989, 80(4), 11-66 (Abstract No. 0263).
 Encyclopedia of Polymer Science and Engineering, vol. 7, Fluorocarbon Elastomers, p. 257-267, Mar. 1989.
 Farb, A. et al., "Vascular smooth muscle cell cytotoxicity and sustained inhibition of neointimal formation by fibroblast growth factor 2-saporin fusion protein," *Circ. Res.*, 1997, 80, 542-550.
 Ferns, G. A. A. et al., "Inhibition of Neointimal Smooth Muscle Accumulation After Angioplasty by an Antibody to PDGF," *Science*, 1991, 253, 1129-1132.
 Fischman, D. L. et al., "A Randomized Comparison of Coronary-Stent Placement and Balloon Angioplasty in the Treatment of Coronary Artery Disease," *N. Eng. J. Med.*, Aug. 25, 1994, 331(8), 496-501.
 Franklin, S. M. et al., "Pharmacologic prevention of restenosis after coronary angioplasty: review of the randomized clinical trials," *Coronary Artery Disease* Mar. 1993, 4(3), 232-242.
 Fukuyama, J. et al., "Tranilast suppresses the vascular intimal hyperplasia after balloon injury in rabbits fed on a high-cholesterol diet," *Eur. J. Pharmacol.*, 1996, 318, 327-332.
 Gregory, C. R. et al., "Rapamycin Inhibits Arterial Intimal Thickening Caused by Both Alloimmune and Mechanical Injury," *Transplantation*, Jun. 1993, 55(6), 1409-1418.

US 7,223,286 B2

Page 6

- Gregory, C. R. et al., "Treatment with Repamycin and Mycophenolic Acid Reduces Arterial Intimal Thickening Produced by Mechanical Injury and Allows Endothelial Replacement," *Transplantation*, Mar. 15, 1995, 59(5), 655-661.
- Guyton, J. R. et al., "Inhibition of rat arterial smooth muscle cell proliferation by heparin. In vivo studies with anticoagulant and nonanticoagulant heparin," *Circ. Res.*, 1980, 46, 625-634.
- Hansson, G. K. et al., "Interferon- γ Inhibits Arterial Stenosis After Injury," *Circ.*, 1991, 84, 1266-1272.
- Hashemolhosseini, S. et al., "Rapamycin Inhibition of the G1 to S Transition Is Mediated by Effects on Cyclin D1 mRNA and Protein Stability," *J Biol Chem*, Jun. 5, 1998, 273, 14424-14429.
- Jonasson, J. et al., "Cyclosporin A inhibits smooth muscle proliferation in the vascular response to injury," *Proc. Natl. Acad. Sci.*, 1988, 85, 2303-2306.
- Lange, R. A. MD et al., "Restenosis After Coronary Balloon Angioplasty," *Annu. Rev. Med.*, 1991, 42, 127-132.
- Liu, M. W. et al., "Trapidil in Preventing Restenosis After Balloon Angioplasty in the Atherosclerotic Rabbit," *Circ.*, 1990, 81, 1089-1093.
- Liu, M. W., MD et al., "Restenosis After Coronary Angioplasty Potential Biologic Determinants and Role of Intimal Hyperplasia," *Circulation*, 1989, 79, 1374-1387.
- Lundergan, C. F. et al., "Peptide inhibition of Myointimal Proliferation by Angiopeptin, a Somatostatin Analogue," *JACC*, May 1991, 17(6), 132B-136B.
- Majesky, M. W. et al., "Heparin regulates smooth muscle S phase entry in the injured rat carotid artery," *Circ. Res.*, 1987, 61, 296-300.
- Marx, S. O. et al., "Rapamycin-FKBP Inhibits Cell Cycle Regulators of Proliferation in Vascular Smooth Muscle Cells," *Circ. Res.*, 1995, 76, 412-417.
- Nemecek, G. M. et al., "Terbinafine Inhibits the Mitogenic Response to Platelet-Derived Growth Factor in Vitro and Neointimal Proliferation in Vivo," *J. Pharmacol. Exp. Thera.*, 1989, 248, 1167-1174.
- Okada, T. et al., "Localized Release of Perivascular Heparin Inhibits Intimal Proliferation after Endothelial Injury without Systemic Anticoagulation," *Neurosurgery*, 1989, 25, 892-898.
- Poon, M. et al., "Rapamycin Inhibits Vascular Smooth Muscle Cell Migration," *J. Clin. Invest.*, Nov. 1996, 98(10), 2277-2283.
- Popma, J. J. et al., "Clinical trials of restenosis after coronary angioplasty," *Circulation*, Sep. 1991, 84(3), 1426-1436.
- Powell, J. S. et al., "Inhibitors of Angiotensin-Converting Enzyme Prevent Myointimal Proliferation After Vascular Injury," *Science*, 1989, 245, 186-188.
- Rensing, B. J. et al., "Coronary restenosis elimination with a sirolimus eluting stent," *European Heart Journal*, 2001, 22, 2125-2130.
- Rodeck, C. et al., "Methods for the Transcervical Collection of Fetal Cells During the First Trimester of Pregnancy," *Prenatal Diagnosis*, 1995, 15, 933-942.
- Ruef, J. MD, et al., "Flavopiridol Inhibits Muscle Cell Proliferation In Vitro and Neointimal Formation In Vivo After Carotid Injury in the Rat," From the Division of Cardiology and Sealy for Molecular Cardiology, University of Texas Medical Branch, Galveston; Accepted Apr. 9, 1999; *Circulation* Aug. 10, 1999, pp. 659-665.
- Serruys, P. W. et al., "A comparison of balloon-expandable-stent implantation with balloon angioplasty in patients with coronary artery disease," *N Engl J Med*, Aug. 25, 1994; 331(8), 489-495.
- Serruys, P. W. et al., "Evaluation of ketanserin in the prevention of restenosis after percutaneous transluminal coronary angioplasty. A multicenter randomized double-blind placebo-controlled trial," *Circulation*. Oct. 1993; 88(4 Pt 1), 1588-1601.
- Serruys, P. W. et al., "Heparin-coated Palmaz-Schatz stents in human coronary arteries. Early outcome of the Benestent-II Pilot Study," *Circulation*, Feb. 1, 1996; 93(3), 412-422.
- Siekierka, J. J., "Probing T-Cell Signal Transduction Pathways with the Immunosuppressive Drugs, FK-506 and Rapamycin," *Immunologic Research*, 1994, 13, 110-116.
- Sigwart, et al., "Intravascular Stents to Prevent Occlusion and Restenosis After Transluminal Angioplasty," *N. Engl. J. Med.*, Mar. 19, 1987, 316, 701-706.
- Simons, M. et al., "Antisense *c-myc* oligonucleotides inhibit intimal arterial smooth muscle cell accumulation in vivo," *Nature*, 1992, 359, 67-70.
- Snow, A. D. et al., "Heparin modulates the composition of the extracellular matrix domain surrounding arterial smooth muscle cells," *Am. J. Pathol.*, 1990, 137, 313-330.
- Sollott, S. J. et al., "Taxol Inhibits Neointimal Smooth Muscle Cell Accumulation after Angioplasty in the Rat," *J. Clin. Invest.*, 1995, 95, 1869-1876.
- van Der Giessen, et al., "Self-expandable Mesh Stents: an Experimental Study Comparing Polymer Coated and Uncoated Wallstent Stents in the Coronary Circulation of Pigs," *Circulation* 1990, 82(suppl. III):III-542.
- van Der Giessen, W. J. et al., "Coronary stenting with polymer-coated and uncoated self-expanding endoprosthesis in pigs," *Coron. Art. Disease* 1992; 3, 631-640.
- Vasey, C. G. et al., "Clinical Cardiology: Stress Echo and Coronary Flow", *Circulation*, Oct. 1989, 80(4) Supplement II, II-66.
- Verweire, E. et al., "Evaluation of Fluorinated Polymers As Coronary Stent Coating," *Journal of Materials Science: Materials in Medicine*, Apr. 2000.
- Weinberger, J. et al., "Intracoronary irradiation: dose response for the prevention of restenosis in swine," *Int. J. Rad. Onc. Biol. Phys.*, 1996, 36, 767-775.
- Preliminary Amendment in U.S. Appl. No. 07/258,189, May 22, 1989.
- Trial Transcript from Nov. 6, 2000 at 185-90 and 235-36 (Attorneys' opening remarks regarding '984 patent).
- Trial Transcript from Nov. 7, 2000 at 274-301, 307-315, 320-28 and 332 (Cordis expert testimony regarding the Palmaz-Schatz stent); 370-379, 480-496 (J. Palmaz testimony regarding the Palmaz-Schatz stent, the '984 patent and the connected z-stent art).
- Trial Transcript from Nov. 8, 2000 at 547-63, 657-63, 674-722, 782-85 (Cordis expert testimony regarding the Palmaz-Schatz stent, the '984 patent and the connected z-stent art).
- Trial Transcript from Nov. 9, 2000 at 819-23, 921 (Cordis expert testimony regarding the '984 patent); 926-941. (R. Croce testimony re Palmaz-Schatz stent); 1033-1053. (R. Schatz testimony).
- Trial Transcript from Nov. 13, 2000 at 1086-1 134. (R. Schatz testimony); 1275-1305 (Cordis expert testimony regarding the '984 patent).
- Trial Transcript from Nov. 14, 2000 at 1390-1404, 1448-1454, 1486-1500 (Cordis expert testimony regarding the '984 patent).
- Trial Transcript from Nov. 15, 2000 at 1686-87, 1724-42, 1828-34, 1850-54, 1887-92 (AVE expert testimony regarding the '984 patent).
- Trial Transcript from Nov. 16, 2000 at 2077-198 (AVE expert testimony regarding the alleged obviousness of the '984 patent).
- Trial Transcript from Nov. 17, 2000 at 2331-34 (jury instructions as to the meaning of the limitations of the claims of the '984 patent).
- Trial Transcript from Nov. 20, 2000 at 2441-48, 2499-2500, 2546-50, 2552-56 (Attorneys' closing arguments regarding the '984 patent).
- Trial Transcript from Nov. 21, 2000 at 2592-94 (reading of jury verdict).
- Trial Transcript from Dec. 18, 2000 at 2750-95 (Cordis expert testimony regarding the Palmaz-Schatz stent during the damages phase).
- Trial Transcript from Dec. 20, 2000 at 3421-88 (AVE expert testimony regarding the Palmaz-Schatz stent during the damages phase).
- Jury verdict, dated Nov. 21, 2000.
- District Court decisions on post-trial motions (194 F. Supp. 2d 323).
- Court of Appeal for the Federal Circuit decision (339 F.3d 1352).
- Trial Transcript from Mar. 4, 2005 at 133-135, 171-173 and 192-96 (Attorney's opening remarks regarding '984 validity).
- Trial Transcript from Mar. 7, 2005 at 275-31 1 (Cordis expert testimony regarding the Palmaz-Schatz stent); 342-46, 353-59, 416-425 (J. Palmaz testimony regarding the Palmaz-Schatz stent, the '984 patent and the connected z-stent art); 430-449, 452-58, 462-492 (R. Croce testimony regarding the Palmaz-Schatz stent); 500-507 (Cordis expert testimony regarding the '984 patent).

US 7,223,286 B2

Page 7

Trial Transcript from Mar. 8, 2005 at 609 (Cordis expert testimony regarding the '984 patent); 628-73, 724-740, 773, 801-839 (Cordis expert testimony regarding the '984 patent), the prior art and the Palmaz-Schatz stent).

Trial Transcript from Mar. 9, 2005 at 936-49, 968-69 (Cordis expert testimony regarding the '984 patent, the prior art and the Palmaz-Schatz stent).

Trial Transcript from Mar. 10, 2005 at 1427-74, 178-1509, 1514-23 (AVE expert testimony regarding the alleged obviousness of the '984 patent); 1566-93 (AVE expert testimony regarding Palmaz-Schatz stent); 1634-49 (R. Schatz testimony).

Trial Transcript from Mar. 11, 2005 at 1846-47, 1891-1900, 1919 (Attorneys' closing arguments regarding '984 obviousness).

Trial Transcript from Mar. 14, 2005 at 1964-67 (reading of jury verdict).

Jury verdict dated Mar. 14, 2005.

Medtronic Vascular Inc.'s Opening Brief in Support of Its Motion for Judgement As A Infringement Claim dated Apr. 19, 2005.

Medtronic Vascular Inc.'s Opening Brief in Support of Its Motion for a New Trial dated Apr. 9, 2005.

D.I. 1407, Cordis' Combined Answering Brief In Opposition to AVE's Motion for JMOL on Infringement of the Palmaz '762 and Schatz '984 Patents and Its Motion for a New Trial dated May 5, 2005.

D.I. 1414, Medtronic Vascular Inc.'s Combined Reply Brief In Support of Its Motion for Judgement as a Matter of Law on Cordis Corp.'s Patent Infringement Claims and Its Motion for a New Trial dated May 19, 2005.

Trial Transcript from Feb. 8, 2001 at 372-412, 449-469 (B. Tobor testimony regarding the prosecution of the '417, '984 and '332 patents); 510-13 (J. Milnamow testimony regarding the prosecution of the '332 patent); 558-604 (J. Palmaz testimony regarding the prosecution of the '417, '984 and '332 patents and the prior art).

Trial Transcript from Feb. 9, 2001 at 637-45, 662-672, 682-85 (J. Palmaz testimony regarding the prior art); 699-742 (R. Schatz testimony); 769-770, 790-95 (Cordis expert testimony regarding prior art).

D.I. 1067, Medtronic AVE, Inc.'s Post-Trial Brief Relating to the Unenforceability of the '762 and '984 Patents Due to Inequitable Conduct.

D.I. 1077, Cordis' Combined Answering Brief in Opposition to AVE's BSC's Post-Hearing Briefs on Alleged Inequitable Conduct Concerning the '762, '984 and '332 Patents.

D.I. 1089, Reply Brief In Support of Medtronic AVE, Inc.'s Contention that the '762 and '984 Patents are Unenforceable Due to Inequitable Conduct dated May 7, 2001.

C.A. No. 00-886-SLR, Answer and Counterclaims of Def. Medtronic AVE, Inc. To First Amended Complaint of Plaintiff Cordis Corp.

BSC's Opening Post-Trial Brief in Support of Its Defense That the Patents in Suit Are Unenforceable, dated Mar. 16, 2001.

Reply Brief in Support of BSC's Defense That the Patents in Suit Are Unenforceable, dated May 7, 2001.

Court's Decision on allegations of inequitable conduct (194 F. Supp. 2d 323) Mar. 28, 2002.

Trial Transcript from Nov. 21, 2000 at 155-57 and 180-84 (Attorneys' opening remarks regarding '332 patent).

Trial Transcript from Nov. 27, 2000 at 227-51, 260-300 (Cordis expert testimony regarding the Palmaz-Schatz stent); 343-60, 363-67, 424-33 (J. Palmaz testimony regarding the Palmaz-Schatz stent and the '332 patent).

Trial Transcript from Nov. 28, 2000 at 649-71.

Trial Transcript from Nov. 29, 2000 at 791-816, 859-870, 953-62 (Cordis expert testimony regarding the '332 patent and the Palmaz-Schatz stent).

Trial Transcript from Nov. 30, 2000 at 1018 (Cordis expert testimony regarding the '332 patent); 1062-80, 1108-1111 (R. Croce testimony regarding the Palmaz-Schatz stent); 1169-70, 1205-17, 1236-45 (Cordis expert testimony regarding the '332 patent).

Trial Transcript from Dec. 1, 2000 at 1352-54 (Cordis expert testimony regarding the '332 patent); 1364-1442 (R. Schatz testimony); 1493-1508, 1552-69 (BSC expert testimony regarding the '332 patent and the Palmaz-Schatz stent).

Trial Transcript from Dec. 4, 2000 at 1602-12, 1638-51, 1713-14, 1730-61, 1811-14, 1823-36 (BSC expert testimony regarding the alleged obviousness of the '332 patent, the prior art and the Palmaz-Schatz stent).

Trial Transcript from Dec. 6, 2000 at 2318-27, 2342-58 (BSC expert testimony regarding the '332 patent).

Trial Transcript from Dec. 7, 2000 at 2549-52 (Cordis expert testimony regarding the '332 patent); 2575-2579, 2591-92, 2630-31, 2649, 2669-71, 2684-85, 2688, 2708-10, 2725-27 (Attorney closing argument regarding '332 patent); 2742-46 Q'ury instructions as to the meaning of the limitations of the claims of the '332 patent).

Trial Transcript from Dec. 11, 2000 at 2817-22 (reading of jury verdict).

Jury verdict, dated Dec. 11, 2000.

D.I. 699, Motion by Defendant BSC and Scimed Life Systems, Inc. For Summary Judgment of Invalidity of U.S. Appl. No. 5,902,332 dated Apr. 4, 2000.

D.I.896, Order Denying Motion for Summary Judgment of Invalidity and Unenforceability of Claims 1, 2, and 5 of the U.S. Appl. No. 5,902,332 Denying {699-1} Motion for Summary Judgment of Invalidity of U.S. Appl. No. 5,902,332 dated Oct. 12, 2000.

Wright et al., Percutaneous Endovascular Stent: An Experimental Study (Abstract), RSNA Meeting (Nov. 28, 1984).

Hearing Transcript from Feb. 10, 1998 at 122-32, 146-80 (Attorneys' opening remarks regarding '417 patent); 180-312 (R. Schatz testimony) [Portions of This Transcript Have Been Removed as Confidential].

Hearing Transcript from Feb. 11, 1998 at 427-575, 577-651 (Cordis expert testimony regarding the '417 patent, the prior art and the Palmaz-Schatz stent).

Hearing Transcript from Feb. 13, 1998 at 1121-1261 (Guidant expert testimony regarding the alleged obviousness of the '417 patent, the prior art and the Palmaz-Schatz stent). [Portions of This Transcript Have Been Removed as Confidential].

Order by J. Robinson denying Cordis' Motion for a Preliminary Injunction Against ACS dated Jul. 17, 1998.

ACS, Inc.'s and Guidant Corp.'s Opening Brief in Support of Their Motion for Summary Judgment of Invalidity of U.S. Appl. No. 5,102,417 dated Aug. 27, 1998.

Plaintiffs's Answering Brief in Opposition to ACS' and BSC's Motion for Summary Judgment on Obviousness dated Sep. 24, 1998.

Order dated Mar. 31, 2000.

Schatz Deposition Testimony; May 15, 1996: 79-83, 89-92, 105-107 and 153-161.

Schatz Deposition Testimony; May 16, 1996: 555-564, 569-572.

Schatz Deposition Testimony; Jan. 8, 1998: 67-73, 108-110.

Schatz Deposition Testimony; Jul. 14, 1998: 69-77, 108-112, 119-123.

Schatz Deposition Testimony; Jul. 12, 1999: 88-91, 132-135, 144-149, 218-223, 231-242.

Schatz Deposition Testimony; Jul. 13, 1999: 251-334, 339-345, 374-416.

Schatz Deposition Testimony; Jul. 14, 1999: 454-550.

Schatz Deposition Testimony; Jul. 15, 1999: 560-614.

Schatz Deposition Testimony; Dec. 2, 1999: 906-911, 928-942, 945-963, 976-978, 1029-1034, 1038-1042.

Palmaz Deposition Testimony, Nov. 5, 1991: 160-172.

Palmaz Deposition Testimony, Feb. 5, 1995: 710-727.

Palmaz Deposition Testimony, Jul. 16, 1998: 55-56; 81-82.

Palmaz Deposition Testimony, Jul. 28, 1999: 560-568, 570-579.

Palmaz Deposition Testimony, Jul. 29, 1999: 778-785.

Palmaz Deposition Testimony, Aug. 31, 1999: 1403-1452.

Palmaz Deposition Testimony, Sep. 2, 1999: 1953-1960.

Palmaz Deposition Testimony, Oct. 14, 1999: 2201-2209; 2275-2342; 2371-2411.

Palmaz Deposition Testimony, Oct. 15, 1999: 2424-2497; 2508-2589.

Palmaz Deposition Testimony, Oct. 16, 1999: 2853-2860.

Tobor Deposition Testimony, Jun. 17, 1999: 837-958.

Tobor Deposition Testimony, Jun. 18, 1999: 1095-1184.

Tobor Deposition Testimony, Dec. 1, 1999: 1217-1371.

US 7,223,286 B2

Page 8

- Tobor Deposition Testimony, Dec. 2, 1999: 1398-1414; 1444-1508; 1532-1548.
- Tobor Deposition Testimony, Dec. 3, 1999: 1652-1653; 1662-1672; 1683-1694.
- Kula Deposition Testimony, Apr. 20, 1999: 268-169.
- Kula Deposition Testimony, Nov. 16, 1999: 660-675; 680-694; 7-8-755; 774-821.
- Kula Deposition Testimony, Nov. 18, 1999; 176-223.
- Expert Report of Dr. Rodney S. Badger on Behalf of Medtronic AVE, Inc. (Jan. 31, 2000).
- Expert Report of Dr. Joseph Bonn on Behalf of Medtronic AVE, Inc. (Jan. 31, 2000).
- Deposition of Dr. Joseph Bonn dated Mar. 14, 2000.
- Rebuttal Expert Report of Nigel Buller, B.Sc., M.B., F.R.C.P. (Mar. 2000).
- Second Supplemental Rebuttal Expert Report of Nigel Buller, B.Sc., M.B., F.R.C.P. (Aug. 17, 2004).
- Rebuttal Expert Report of John M. Collins, PH.D. (Feb. 2000).
- Expert Report of David C. Cumberland, M.D. (Jan. 24, 2000).
- Expert Report of John T. Goolkasian (Feb. 2000).
- Deposition of Richard R. Heuser, M.D. (Sep. 7, 2004).
- Deposition of Henry R. Piehler (Sep. 10, 2004).
- Deposition of Ronald J. Solar (Mar. 22, 2000).
- Deposition of Ronald J. Solar (Mar. 23, 2000).
- Deposition of Ronald J. Solar (Apr. 12, 2000).
- Expert Report of Dr. Arina Van Breda on Behalf of Medtronic AVE, Inc. (Jan. 31, 2000).
- Deposition of Anna Van Breda (Mar. 24, 2000).
- Deposition of Arina Van Breda (Aug. 21, 2004).
- Expert Report of John F. Witherspoon (Jan. 24, 2000).
- Supplemental Expert Report of John F. Witherspoon (Oct. 27, 2000).
- Deposition of John F. Witherspoon (Mar. 8, 2000).
- Palmaz et al., Article: "Normal and Stenotic Renal Arteries: Experimental Balloon Expandable Intraluminal Stenting", Radiology, Sep. 1987. (AVE 84).
- Julio C. Palmaz, Article: "Expandable vascular endoprosthesis." (AVE 132).
- Duprat et al., Article: Flexible Balloon-Expandable Stent for Small Vessels Duprat et al. Radiology, vol. 162, pp. 276-278, 1987. (AVE 134).
- Coons et al., Article: "Large-Bore, Long Biliary Endoprosthesis (Biliary Stents) for Improved Drainage," Radiology, vol. 148, pp. 89-94, 1983. (AVE 143).
- Honickman et al., Article: "Malpositioned Biliary Endoprosthesis, Technical Developments And Instrumentation," vol. 144, No. 2., 1982. (AVE 144).
- Harries-Jones, et al., Article: "Repositioning of Biliary Endoprosthesis with Gruntzig Balloon Catheters," AJR, vol. 138, pp. 771-772, 1982. (AVE 153).
- Charnsangavej et al., Article "Stenosis of the Vena Cava: Preliminary Assessment of Treatment with Expandable Metallic Stents," Radiology, vol. 161, pp. 295-298, 1986. (AVE 359).
- Wallace, M. J. et al., Article "Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications," Radiology, vol. 158, pp. 309-312, 1986. (AVE 364).
- T. Yoshioka, et al., AIR Article: "Self-Expanding Endovascular Graft: An Experimental Study in Dogs", vol. 151, pp. 673-676, 1988. (AVE 438).
- Palmaz, J. C. et al., Article: "Expandable Intraluminal Vascular Graft: A Feasibility Study," Surgery, vol. 99, pp. 199-205, 1986. (AVE 461).
- Lawrence et al., Article: "Percutaneous Endovascular Graft: Experimental Evaluation." Radiology, vol. 163, pp. 357-360, 1987. (AVE 671).
- Palmaz et al., Article: Expandable Intraluminal Graft: A Preliminary Study, Nov. 17-22, 1985, Radiology, vol. 156, pp. 73-77, 1985. (AVE 1224).
- Fallone et al., "Elastic Characteristics of the Self-Expanding Metallic Stents," Investigative Radiology, vol. 23, pp. 370-376, 1988. (AVE 1953).
- Palmaz Paper Entitled "Research Project Expandable Vascular Endoprosthesis" May 18, 1983.
- Rousseau, et al., Publication: "Percutaneous Vascular Stent: Experimental Studies & Preliminary Clinical Results in Peripheral Arterial Diseases," in Inter. Angio, vol. 6, 153-161, 1987. (AVE 3301).
- Rousseau, et al., Publication: "Self-Expanding Endovascular Prosthesis: An Experimental Study," Radiology, vol. 164, pp. 709-714, 1987. (AVE 3303).
- Wallace, et al., Article: "Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications," Radiology, vol. 58, pp. 309-312, 1986. (DBX 2938).
- Palmaz et al., Article: "Expandable Intraluminal Graft: A Preliminary Study," Radiology, vol. 156, pp. 73-77, Nov. 17-22, 1985 (DBX 4595).
- Program for the 12th Annual Course on Diagnostic Angiography and Interventional Radiology Mar. 23-26, 1987 sponsored by The Society of Cardiovascular and Interventional Radiology (DBX 6235).
- Preliminary Motion for Judgment re: Wolff claims 1, 2-8, 10, 15 and 19 (DBX6759).
- Palmaz Declaration (DBX 7069).
- Letter from Gaterud to Dr. Palmaz dated Jul. 5, 1988 with attached document entitled: "Segmented, balloon-expandable stents." (DBX 7160).
- Duprat et al., Article: "Flexible Balloon-Expandable Stent For Small Vessels," Radiology, vol. 168, pp. 276-278, 1987 (PX 82).
- Drawing Sent to Bodic on Mar. 17, 1986 (PX 374).
- Letter from Dr. Palmaz to R. Bowman enclosing a model of the flexible coronary graft dated Mar. 17, 1986 (PX 337).
- Lab Notebook pages dated Jul. 30, 1987 from Rodney Wolff (COR 185596-597) (PX621A).
- Charnsangavej, et al., Article: "Stenosis of The Vena Cava Preliminary Assessment of Treatment with expandable Metallic Stents," Radiology, vol. 161, No. 2, pp. 295-298 with attached photographs, 1986. (API 72).
- J. Palmaz: The Current Status of Vascular Prosthesis, published by SCIR in the Twelfth Annual Course on Diagnostic Angiography And Interventional Radiology Mar. 23-26, 1987. (API 73).
- Amendment in Response to Office Action of Oct. 18, 1998 in re: Application of Julio Palmaz U.S. Appl. No. 174,246. (API 152).
- Article: Wallace, et al., Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications Work In Progress, Radiology, vol. 158, pp. 309-312. (API 295).
- Reply of Senior Party Schatz To Patentee Wolffs Opposition To The Belated Motion For Judgment Of Applicant Schatz With Regard To Wolff Claims 1, 2-8, 10, 11, 13-17, And 19 (COR 186450-455) (API 310).
- Brief Of Senior Party Schatz At Final Hearing (API 313).
- Copy of Letter from Ron Sickles to Ben Tobor dated Feb. 10, 1988 (Exhibit 42).
- Copy of Letter from R.O. Sickles to Mike Tatlow dated May 12, 1988 (Exhibit 43).
- Copy of Letter from R. O. Sickles to Richard Schatz dated Jun. 2, 1988 (Exhibit 44).
- Copy of Letter from Richard Schatz to Raimund Erbel dated Jun. 3, 1988 (Exhibit 45).
- Copy of Letter from Richard Schatz to Mike Schuler dated Aug. 29, 1991 (Exhibit 48).
- Minutes of J&J Stent Project Review Meeting datd Jan. 21, 1988 (Exhibit 7).
- Preliminary Motion for Judgment with Regard to Wolff Claims 1, 2-8, 10, 11, 13-17, and 19. (Exhibit 67).
- Declaration of Richard A Schatz. (Exhibit 75).
- Belated Motion for Judgement with Regard to Wolff Claims 1, 2-8, 10, 11, 13-17 and 19. (Schatz—Exhibit 77).
- Letter from Dr. Schatz to Mr. Tobor, dated Jun. 3, 1988. (Exhibit 122).
- Letter from Dr. Schatz to Mr. Romano, dated Nov. 28, 1988. (Exhibit 131).
- Letter from Mr. Sickles to Mr. Tobor, dated Feb. 10, 1988 (Exhibit 145).
- Richard A. Schatz, Article title: "A View of Vascular Stents" Circulation, vol. 79, No. 2, pp. 445-457, 1989. (Exhibit 194).

US 7,223,286 B2

Page 9

- Senior Party Schatz's reply to Patentee Wolffs Opposition to the Preliminary Motion Of Application Schatz for judgment with regard to Wolff Claims 1, 2-8, 10, 11, and 13-17. (Exhibit 69).
- Wallace, et al., Article: "Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications' Work In Progress," Radiology, vol. 158, pp. 309-312, 1986. (Exhibit 165).
- Charnsangavej, et al., Article: "Stenosis of The Vena Cava Preliminary Assessment of Treatment with expandable Metallic Stents," Radiology, vol. 161, No. 2, pp. 295-298 with attached photographs, 1986! (Exhibit 167).
- David D. Lawrence et al., Publication: Percutaneous Endovascular Graft: Experimental Evaluation¹, Radiology, pp. 163, 357-360, 1987. (Exhibit 173).
- Charles E. Putnam, M.D., Cover and article from "Investigative Radiology", vol. 23, No. 5, May 1988. (Exhibit 177).
- Robert N. Berk, Cover and article from "American Journal of Roentology", pp. 673-676, 1988. (Exhibit 178).
- Declaration of John S. Kula Under 37 CFR § 1.672. (Kula—Exhibit 77).
- Yoshioka et al., Article: "Self-Expanding Endovascular Graft: An Experimental Study in Dogs" AJR, vol. 151, pp. 673-676, 1988. (PX 100).
- Palmaz, et al., Article: Expandable Intraluminal Graft: A Preliminary Study Work in Progress¹, Radiology, vol. 156, No. 1, pp. 73-77, 1985. (PX 101).
- Declaration of Richard Schatz Under 37 C.F.R. § 1.672. (PX 106).
- Charnsangavej et al., Article: "Stenosis of the Vena Cave: Preliminary Assessment of Treatment with Expandable Metallic Stents," Radiology, vol. 161, pp. 295-298, 1986. (PX 143).
- Wallace, et al., Article: Tracheobronchial Tree: Expandable Metallic Stents Used in Experimental and Clinical Applications Work in Progress¹, Radiology, vol. 158, pp. 309-312, 1986. (PX 144).
- Gina Kolata, News Article: NY Times, "Devices That Opens Clogged Arteries Gets a Falling Grade in a New Study", pp. 16-18, Jan. 3, 1991. (PX 186).
- Duprat, et al., Article: "Flexible Balloon-Expanded Stent for Small Vessels Work in Progress¹", Radiology, vol. 162, pp. 276-278, 1987. (PX 207).
- Letter from Palmaz to Bowman dated Mar. 17, 1986. (PX 350).
- Memo re: Minutes of Stent Project Review- San Antonio- Mar. 15, 1988. (PX 651).
- Kuntz, et al., Article: Clinical Cardiology Frontiers: "Defining Coronary Restenosis, Newer Clinical and Angiographic Paradigms", Circulation, Sep. 1993, vol. 88, No. 3, pp. 1310-1323. (PX 854).
- Belated Motion for Judgment with regard to Wolff Claims 1, 2-8, 10, 11, 13-17, and 19. (PX 1410).
- Drawing of Spiral Stent (sent to Bodic Mar. 17, 1986). (PX2933).
- Wright et al., Article: "Percutaneous Endovascular Stents: An Experimental Evaluation," Radiology, vol. 156, pp. 69-72, 1985. (PX 3093).
- Charnsangavej et al., Article: "A New Expandable Metallic Stent for Dilation of Stenotic Tubular Structures: Experimental and Clinical Evaluation," Houston Medical Journal, vol. 3, pp. 41-51, Jun. 1987. (PX 3207).
- In re Application of Wiktor, U.S. Appl. No. 69,636, Response to Office Action dated Mar. 17, 1988. (PX3236).
- Transmittal Letter of Response to First Office Action in '417 patent. (PX 3993).
- Letter from B. Tobor to R. Schatz dated Jul. 23, 1991. (PX 3996).
- Mullins et al., Article: "Implantation of balloon-expandable intravascular grafts by catheterization in pulmonary arteries and systemic veins," Circulation, vol. 77, No. 1, pp. 188-189, 1988. (PX4049).
- Schatz et al., Article: "Intravascular Stents for Angioplasty," Cardio, 1997. (PX 4050).
- Schatz et al., Article: "New Technology in Angioplasty Balloon-Expandable Intravascular Stents, New Developments in Medicine," vol. 2, No. 2, pp. 59-75, 1987. (PX4051).
- Richard A. Schatz, Article: "Introduction to Intravascular Stents," Cardiology Clinics, vol. 6, No. 3, pp. 357-372, 1988. (PX 4052).
- Richard A. Schatz, Article: "A View of Vascular Stents," Circulation, vol. 79, No. 2, pp. 445-457, 1989. (PX4053).
- Wang et al., Article: "An Update on Coronary Stents," Cardio, pp. 177-186, 1992. (PX 4054).
- Richard A. Schatz, Article: "New Technology in Angioplasty: Balloon-Expandable Starts," Medicamundi, vol. 33, No. 3, pp. 1121-1116, 1988. (PX 4055).
- Letter from Tobor to Schatz dated Sep. 29, 1988. (PX 1395).
- Verified Statement of Facts by Innamed Inventor R.A. Schatz document filed in U.S. Patent and Tradement Office on Sep. 8, 1989. (PX 3677).
- Declaration of John S. Kula Under 37 CFR § 1.672 (Exhibit 329).
- Letter to Mike Schular from R.A. Schatz dated Aug. 29, 1991. (Exhibit 402).
- Articulated, Balloon—Expandable Stents, (DBX 7159).
- J. Rosch et al., Experimental Intrahepatic Portacaval Anastomosis: Use of Expandable Gianturco Stents, Radiology, vol. 162, pp. 481-485, 1987.
- J. Rosch et al., Modified Gianturco Expandable Wire Stents In Experimental and Clinical Use, Ann Radiol, vol. 31, No. 2, pp. 100-103, 1987.
- J. Rosch et al., Gianturco Expandable Stents In the Treatment of Superior Vena Cava Syndrome Recurring After Vena Cava Syndrome Recurring After Maximum-Tolerance Radiation, Cancer, vol. 60, pp. 1243-1246, 1987.
- I.E. Gordon, Structures or Why Things Don't Fall Down, Penguin Books, pp. 45-59, 132-148, 210-244, 377-383.
- Maass et al., Radiological Follow-up of Transluminally Inserted Vascular Endoprostheses: An Experimental Study Using Expanding Spirals, Radiology, vol. 152, pp. 659-663, 1984.
- Argument submitted re EP 861 15473 dated Jan. 20, 1995. (AVE 2478).
- Verified Statement of Facts by Julio C. Palmaz dated Aug. 4, 1989. (PX 3662).
- Papanicolaou et al., Insertion of a Biliary Endoprosthesis Using A Balloon Dilatation Catheter, Gastrointest Radiology, vol. 10, pp. 394-396, 1985.
- Palmaz et al., Atherosclerotic Rabbit Aortas: Expandable Intraluminal Grafting, Radiology, vol. 168, pp. 723-726, 1986.
- Palmaz, The Current Status of Vascular Prostheses; Rosch et al., Gianturco, Expandable Stents in Experimental and Clinical Use, SCIVR, pp. 118-124, 1987.
- Rosch et al., Abstract: Modified Gianturco Expandable Wire Stents in Experimental and Clinical Use, CIRSE, Porto Cervo, Sardinia, May 25-29, 1987.
- Rosch et al., Gianturco Expandable Wire Stents in the Treatment of Superior Vena Cava Syndrome Recurring After Maximum-Tolerance Radiation, Cancer, vol. 60, pp. 1243-1246, 1987.
- Mirich et al., Percutaneously Placed Endovascular Grafts for Aortic Aneurysms: Feasibility Study, Radiology, vol. 170, pp. 1033-1037, 1989.
- Dotter, Transluminally-placed Coilspring Endarterial Tube Grafts, Investigative Radiology, vol. 4, Sep.-Oct., pp. 329-332, 1969.
- Palmaz et al., Abstract: Expandable Intraluminal Graft: A Preliminary Study, Radiology, vol. 153 (P), Nov. 1983: 70th Scientific Assembly and Annual Meeting.
- Cragg et al, Nonsurgical Placement of Arterial Endoprosthesis: A New Technique Using Nitinol Wire, Radiology, vol. 147, pp. 261-263, Apr. 1983.
- J. Rosch et al., Gianturco Expandable Stents in Experimental and Clinical Use, Program: "Twelfth Annual Course on Diagnostic Angiography and Interventional Radiology" (Society of Cardiovascular and Interventional Radiology, Pittsburgh, PA), Mar. 23-26, 1987 (the second Monofilament Article).
- Uchida et al., Modifications of Gianturco Expandable Wire Stents, AIR, vol. 150, pp. 1185-1187, 1988.
- Palmaz, Balloon-Expandable Intravascular Stent, AJR, vol. 1510, pp. 1263-1269.
- Cordis Corporation v. Advanced Cardiovascular Systems, Inc., Guidant Corporation, Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCMED Life Systems, Inc., Plaintiffs Complaint, Oct. 23, 1997 (Case No. 97-550-SLR).
- Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership, Plaintiffs First Amended Complaint for Declaratory Relief of Patent Validity,

US 7,223,286 B2

Page 10

Unenforceability, Noninfringement, and for Antitrust Violations, Jan. 27, 1998 (Civil Action No. 97-700).

Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership, Cordis Corporation and Johnson & Johnson's Answer and Counterclaim, Feb. 27, 1998 (Civil Action No. 97-700-SLR).

Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership, Expandable-Graft Partnership's Answer, Feb. 27, 1998 (Civil Action No. 97-700-SLR).

Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership, Reply of Plaintiff Arterial Vascular Engineering, Inc. To Counterclaims of Defendant Cordis Corporation, Mar. 31, 1998 (Civil Action No. 97-700-SLR).

Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable-Grafts Partnership, Reply of Plaintiff Arterial Vascular Engineering, Inc. To Counterclaims of Defendant Expandable Grafts Partnership, Mar. 31, 1998 (Civil Action No. 97-700-SLR).

Cordis Corporation v. Advanced Cardiovascular Systems, Inc. and Guidant Corporation, Cordis Corporation's Motion for a Preliminary Injunction, Oct. 8, 1997 (Civil Action No. 97-550).

Cordis Corporation v. Advanced Cardiovascular Systems, Inc., Guidant Corporation Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCJVID, Inc., Cordis's Motion for Preliminary Injunction Against Arterial Vascular Engineering, Inc., Dec. 29, 1997 (Case No. 97-550-SLR).

Deposition of R. Schatz, M.D. in Cordis Corporation v. Advanced Cardiovascular Systems, Inc., taken on Jan. 8, 1998 (Civil Action No. 97-550 SLR).

Deposition of Lee P. Bendel in Cordis Corporation v. Advanced Cardiovascular Systems, Inc., taken on Jan. 22, 1998 (Civil Action No. 97-550 SLR).

Deposition of Julio Cesar Palmaz in Cordis Corporation v. Advanced Cardiovascular Systems, Inc., taken on Dec. 29, 1997 (Civil Action No. 97-550 SLR).

Deposition of Richard A. Bowman in Cordis Corporation v. Advanced Cardiovascular Systems, Inc., taken on Jan. 9, 1998 (Civil Action No. 97-550 SLR).

Deposition of Gary Schneiderman in Cordis Corporation v. Advanced Cardiovascular Systems, Inc., taken on Jan. 16, 1998 (Civil Action No. 97-550 SLR).

Deposition of David Pearle, M.D. in Cordis Corporation v. Advanced Cardiovascular Systems, Inc., taken on Jul. 10, 1998 (Civil Action No. 97-550 SLR).

Preliminary Injunction hearing testimony taken on Feb. 9-13, 1998 (Civil Action No. 97-550 SLR).

Cordis Corporation v. Advanced Cardiovascular Systems, Inc., et al., (Civil Action No. 97-550 SLR) and Cordis Corporation v. Advanced Cardiovascular Systems, Inc. Et al. (Civil Action No. 98-65-SLR), Opening Post Hearing Brief of Plaintiff Cordis Corporation in Support of Motion for Preliminary Injunction, Mar. 6, 1998 (Portions relevant to patent claim construction and patent validity issues).

Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc. et al., Post-Hearing Reply Brief of Plaintiff Cordis Corporation in Support of Its Motion for Preliminary Injunction, Apr. 10, 1998 (Case No. 97-550 SLR) (Portions relevant to patent validity issues).

Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc. et al., Plaintiffs Motion for a Preliminary Injunction Against Boston Scientific Corporation and SCLMED Life Systems, Inc. And Memorandum in Support, Apr. 13, 1998 (Case No. 97-550-SLR).

Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc., et al., Judge Robinson's Order Denying Plaintiffs Motion for a Preliminary Injunction, Jul. 17, 1998 (Civil Action No. 97-550 SLR).

Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc., et al., Defendant Boston Scientific Corporation and SCTMED Life Systems, Inc.'s Motion for Summary Judgment of Invalidity of U.S. Appl. No. 5,102,417, Aug. 27, 1998 (Civil Action No. 97-550-SLR).

Boston Scientific Limited, et al. v. Expandable Grafts Partnership, Plaintiffs' Statement of Claim, Mar. 13, 1997 (UK Action No. 1493).

Boston Scientific Limited, et al. v. Expandable Grafts Partnership, Defendant's Amended Defense and Counterclaim, Aug. 14, 1997 (UK Action No. 1493).

Boston Scientific Limited, et al. v. Expandable Grafts Partnership, Petition for Revocation, Mar. 13, 1997 (UK Action No. 1497).

Boston Scientific Limited, et al. v. Expandable Grafts Partnership, Particulars of Objections, Mar. 13, 1997 (UK Action No. 1497).

Boston Scientific Limited, et al. v. Expandable Grafts Partnership and Boston Scientific Limited et al., v. Julio C. Palmaz, Boston's Skeleton Argument (UK Action Nos. 1493, 1495, 1496, and 1497).

Boston Scientific Limited, et al. v. Julio C. Palmaz and Expandable Grafts Partnership, Skeleton Argument of Palmaz/EGP, Mar. 19, 1998 (UK Action Nos. 1493, 1495, 1496 and 1497).

Boston Scientific Limited, et al. v. Julio C. Palmaz and Expandable Grafts Partnership, EGP's Final Submissions, Apr. 2, 1998 (UK Action Nos. 1493, 1495, 1496 and 1497).

Boston Scientific Limited, et al. v. Julio C. Palmaz and Expandable Grafts Partnership, Judgment, Jun. 26, 1998 (UK Action Nos. 1493, 1495, 1496 and 1497).

Rosch, Modified Gianturco Expandable Wire Stents in Experimental and Clinical Use, CJJR.SE 1987 Presentation: see Witness Statement of Josef Rosch from U.K. Proceeding.

Statement of Claim by Boston Scientific et al. against Expandable Grafts Partnership et al., in EPG et al., v. Boston Scientific et al. in Netherlands (Mar. 13, 1997).

Motion for Joinder of Actions, Change of Claim and Statement of Claim filed by Expandable Grafts Partnership et al. in EPG et al. v. Boston Scientific et al. In Netherlands (Apr. 22, 1997).

Opinion of K.J. Merman filed in EPG et al. v. Boston Scientific et al. in Netherlands (Aug. 29, 1997).

Expert report of Dr. Nigel Buller in EPG et al. v. Boston Scientific et al. in Netherlands (Aug. 28, 1997).

Expert report of Lee P. Bendel in EPG et al. v. Boston Scientific et al. in Netherlands (Aug. 28, 1997).

Memorandum of Oral Pleading in EPG et al. v. Boston Scientific et al. in Netherlands (Sep. 12, 1997).

Plea Notes of P. A.M. in EPG et al. v. Boston Scientific et al. in Netherlands (Mar. 10, 1998).

Decision of Court of Appeals in EPG et al. v. Boston Scientific et al. in Netherlands (Apr. 23, 1998).

Translation of Nullity Action Against EPO 0 364 787 by Biotronik in Germany.

Translation of Nullity Action Against EPO 0 335 341 by Biotronik in Germany.

Translation of EPG Response to Nullity Action Against EP 0 364 787 by Biotronik in Germany.

Translation of EPG Response to Nullity Action EP 0 335 341 by Biotronik in Germany.

Nullity Suit Against EP-B1-0 335 341 Brought by Boston Scientific in Germany.

Translation of Opposition filed by Terumo Corp. Against Japan Patent No. 2680901.

Translation of Decision on Opposition Against Japan Patent No. 2680901.

Memorandum Order of the Court dated Sep. 7, 2000, concerning disputed claim construction.

Translation of Judgment in Nullity Action Against EP 0 364 787 by Biotronik in Germany.

Translation of Judgment in Nullity Action Against EP 0 335 341 by Biotronik in Germany.

Trial transcript from Mar. 17, 2005 at 171-172, 191-192.

Trial transcript from Mar. 18, 2005 at 282-285, 325-327, 349-351.

Trial transcript from Mar. 21, 2005 at 721-726.

Trial transcript from Mar. 24, 2005 at 1387.

Trial transcript from Jul. 26, 2005.

BSC's Opening Brief in Support of Its Motion for Judgment as a Matter of Law or, in the Alternative, for a New Trial, dated Mar. 16, 2001.

US 7,223,286 B2

Page 11

Cordis' Answering Brief in Opposition to BSC's Motion for JMOL or a New Trial on the Palmaz '762 Patent and the Schatz '332 Patents, dated Apr. 17, 2001.

BSC's Reply Brief in Support of Its Motion for Judgment as a Matter of Law or, in the Alternative, for a New Trial, dated May 11, 2001.

J. Rosch et al., Abstract, Expandable Gianturco-Type Wire Stents in Experimental Intrahepatic Portacaval Shunts, Program: "72nd Scientific Assembly and Annual Meeting of the Radiological Society of North America", Nov. 30-Dec. 5, 1986m Radiology, vol. 161, pp. 40-41, 1986.

Cordis Corporation v. Boston Scientific, Order Dated Mar. 27, 2006 (97-550-SLR).

Cordis Corporation v. Boston Scientific, Judgment in a Civil Case Dated Mar. 27, 2006 (97-550-SLR).

Cordis Corporation v. Boston Scientific, Memorandum Opinion Dated Mar. 27, 2006 (97-550-SLR).

Cordis Corporation and Expandable Grafts Partnership v. Advanced Cardiovascular Systems, Inc., Guidant Corporation, Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCIMED Life Systems, Inc., Answer and Counterclaims of Defendant Advanced Cardiovascular Systems, Inc., Apr. 8, 1998 (Case No. 97-550-SLR).

Boston Scientific Limited et al. v. Expandable Grafts Partnership and Boston Scientific Limited et al. v. Julio C. Palmaz, Boston's Closing Submissions (UK Action Nos. 1493, 1495, 1496 and 1497). *Cordis Corporation v. Advanced Cardiovascular Systems, Inc., Guidant Corporation, Arterial Vascular Engineering, Inc., Boston Scientific Corporation and SCIMED Life Systems, Inc.*, Defendants' Answer, Nov. 12, 1997 (Case No. 97-550-SLR).

Statement of Rejoinder in the Action on the Merits, Also Including an Amendment of Defendant's Final Position in the Principal Action, as Well as the Provisional Statement of Rejoinder in the Action on the Counterclaim in *EPG et al. v. Boston Scientific et al.* in Netherlands (Feb. 10, 1998).

Statement of Answer in the Ancillary Appeal in *EPG et al. v. Boston Scientific et al.* in Netherlands (Mar. 10, 1998).

Appeal filed by Expandable Grafts Partnership et al. in *EPG et al. v. Boston Scientific et al.* in Netherlands (Nov. 12, 1997).

Title filed by Boston Scientific et al. in *EPG et al. v. Boston Scientific et al.* in Netherlands (Jan. 22, 1998).

Deposition of Richard Schatz, M.D. in *Cordis Corporation v. Advanced Cardiovascular Systems, Inc.* taken on Jul. 14, 1998 (Civil Action No. 97-550-SLR).

Jury Verdict form from the *Cordis Corporation et al. v. Boston Scientific Corporation, et al.* liability trial, undated.

Trial testimony transcripts from the *Cordis Corporation et al. v. Boston Scientific Corporation et al.* liability trial dated Nov. 21, Nov. 27-Dec. 1, Dec. 4-8 and Dec. 11, 2000.

Boston Scientific SCIMED, Inc. and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc., Opening Expert Report of Stephen R. Hanson, Ph.D. (Civil Action No. 03-328-SLR).

Boston Scientific SCIMED, Inc. and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc., Opening Expert Report of Robson F. Storey, Ph.D. (Civil Action No. 03-283-SLR).

Boston Scientific SCIMED, Inc. and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc., Rebuttal Expert Report of Kinam Park, Ph.D. (Civil Action No. 03-283-SLR).

Cordis Corporation v. Boston Scientific Corporation and SCIMED Life Systems, Inc. (C.A. No. 03-027-SLR) and *Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.* (C.A. No. 03-283-SLR) Combined Post-Hearing Brief In Support Of Cordis Corporation's Motion For Preliminary Injunction in C.A. No. 03-027-SLR, And In Opposition to Plaintiffs' Motion For Preliminary Injunction in C.A. No. 03-283-SLR.

Cordis Corporation v. Boston Scientific Corporation and SCIMED Life Systems, Inc. (C.A. No. 03-027-SLR) *Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Cor-*

poration and Johnson and Johnson, Inc. (C.A. No. 03-283-SLR), Boston Scientific's Opening Post-Hearing Brief.

Cordis Corporation v. Boston Scientific Corporation and SCIMED Life Systems, Inc. (C.A. No. 03-027-SLR) *Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc.* (C.A. No. 03-283-SLR), Combined Post-Hearing Answering Brief In Support of Cordis Corporation's Motion For Preliminary Injunction In C.A. No. 03-027-SLR, And In Opposition To Plaintiffs' Motion For Preliminary Injunction in C.A. No. 03-283-SLR.

Wu et al., Silicone-covered self-expanding metallic stents for the palliation of malignant esophageal obstruction and esophagorespiratory fistulas: experience in 32 patients and a review of the literature, *Gastrointestinal Endoscopy*, 1994, pp. 22-33, vol. 40, No. 1, Portland Oregon.

Binmoeller, et al., Silicone-Covered Expandable Metallic Stents in the Esophagus: An Experimental Study, *Endoscopy*, 1992, pp. 416-420, vol. 24, Georg Thieme Verlag Stuttgart New York.

Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc., Answering Memorandum in Opposition to Plaintiffs Motion for a Preliminary Injunction and Appendix thereto (Civil Action No. 03-283-SLR).

Boston Scientific SCIMED, Inc., and Boston Scientific Corporation v. Cordis Corporation and Johnson and Johnson, Inc., Answering Memorandum in Opposition to Plaintiffs Motion for a Preliminary Injunction and Appendix thereto (Civil Action No. 03-283-SLR).

Rhine, Polymers for Sustained Macromolecule Release: Procedures to Fabricate Reproducible Delivery Systems and Control Release Kinetics, *Journal of Pharmaceutical Sciences*, 1 980, pp. 265-270, vol. 69, No. 3.

Langer et al., Controlled Release of Macromolecules From Polymers, *Biomedical Polymers Polymeric Materials and Pharmaceuticals for Biomedical Use*, 1980, pp. 112-137, Academic Press, Inc., New York, NY.

Langer et al., Applications of Polymeric Delivery Systems for Macromolecules and Factors Controlling Release Kinetics.

Rhine et al., A Method to Achieve Zero-Order Release Kinetics From Polymer Matrix Drug Delivery Systems, pp. 67-72.

Langer et al., Polymers for the Sustained Release of Macromolecules: Controlled and Magnetically Modulated Systems, *Better Therapy With Existing Drugs: New Uses and Delivery Systems*; 1981, pp. 179-216, Merck Sharp & Dohme International, Rahway, NJ.

Hsieh, et al., Zero-Order Controlled-Release Polymer Matrices for Micro-and-Macromolecules, *Journal of Pharmaceutical Sciences*, 1983 pp. 17-22, vol. 72, No. 1.

Brown et al., In Vivo and In Vitro Release of Macromolecules from Polymeric Drug Delivery Systems, *Journal of Pharmaceutical Sciences*, 1983, pp. 1181-1185, vol. 72, No. 10.

Langer, Implantable Controlled Release Systems, *Pharmac. Ther.*, 1983, pp. 35-51, vol. 21, printed in Great Britain.

Kost et al., Controlled Release of Bioactive Agents, *Trends in Biotechnology*, 1984, pp. 47-51, vol. 2, No. 2, Elsevier BV Amsterdam.

Bawa et al., An Explanation for the Controlled Release of Macromolecules from Polymers, *Journal of Controlled Release*, 1985, pp. 259-267, vol. 1 Elsevier Science BV Amsterdam.

Leong et al., Polymeric controlled drug delivery, 1987, pp. 199-233, vol. 1/3, Elsevier Science Publishers BV Amsterdam.

Langer, Polymeric Delivery Systems, *Targeting of Drugs 2 Optimization Strategies*, 1989, pp. 165-174, Plenum Press, New York and London.

Langer, Biomaterials in Controlled Drug Delivery: New Perspective from Biotechnological Advances; *Pharmaceutical Technology*, 1989, pp. 18, 23-24, 26, 28, 30.

Langer, Controlled Release Systems, pp. 115-124.

Laurencin et al., Polymeric Controlled Release Systems: New Methods for Drug Delivery, *Clinics in Laboratory Medicine*, 1987, pp. 301-323, vol. 7, No. 2, WB Saunders Company, Philadelphia.

Langer, Biopolymers in Controlled Release Systems, *Polymeric Biomaterials*, pp. 161-169.

US 7,223,286 B2

Page 12

- Tsong-Pin Hsu et al., Polymers for the Controlled Release of Macromolecules: Effect of Molecular Weight of Ethylene-vinyl Acetate Copolymer, *Journal of Biomedical Materials Research*, 1985, pp. 445-460, vol. 19.
- Langer, Polymers and Drug Delivery Systems, *Long-Acting Contraceptive Delivery Systems*, 1983, pp. 23-32, Harper & Row, Philadelphia, PA.
- Langer, New Drug Delivery Systems: What the Clinician Can Expect, *Drug Therapy*, 1983, pp. 217-231.
- Langer, et al., Chemical and Physical Structure of Polymers as Carriers for Controlled Release of Bioactive Agents: A Review, *Rev. Macromol. Chem. Phys.*, 1983, pp. 61-126.
- Langer, Polymeric Delivery Systems for Controlled Drug Release, *Chem. Eng. Commun.* 1980, pp. 1-48-vol. 6, Gordon and Breach Science Publishers, Inc. USA.
- Langer, et al., Biocompatibility of Polymer Delivery Systems for Macromolecules, *Journal of Biomedical Materials Research*, 1981, pp. 267-277, vol. 15.
- Langer, Controlled Release: A New Approach to Drug Delivery, *Technology Review*, 1981, pp. 26-34.
- Langer, et al., Sustained Release of Macromolecules from Polymers, *Polymeric Delivery Systems*, PGS. 175-176, Gordon and Breach Science Publishers, New York.
- Langer, Polymers for the Sustained Release of Proteins and other Macromolecules, *Nature*, 1976, pp. 797, 263, 799-800, vol. 263, No. 5580.
- Baker, et al., Controlled Release: Mechanisms and Rates (1974).
- Hanson, et al., In Vivo Evaluation of Artificial Surfaces with a Nonhuman Primate Model of Arterial Thrombosis, *Lab Clin. Med.*, Feb. 1980, pp. 289-304.
- Baker, Controlled Release of Biologically Active Agents (1987) pp. 1-275.
- Cordis Corporation v. Boston Scientific Corporation* (CA. No. 03-27-SLR) and *Boston Scientific Scimed, Inc., v. Cordis Corporation and Johnson & Johnson, Incorporated* (CA. No. 03-283-SLR) Hearing Transcripts for Jul. 21, 2003, Jul. 22, 2003, Jul. 23, 2003.
- Cordis Corporation v. Boston Scientific Corporation et al.* (CA. No. 03-027-SLR), and *Boston Scientific Scimed, Inc. et al. v. Cordis Corporation et al.* (CA. No. 03-283-SLR), Boston Scientific's Post-Hearing Reply Brief and Exhibits Thereto, Sep. 12, 2003.
- Cordis Corporation v. Boston Scientific Corporation et al.* (CA. No. 03-027-SLR), and *Boston Scientific Scimed, Inc. et al. v. Cordis Corporation et al.* (CA. 03-283-SLR), Memorandum Order, Nov. 21, 2003.
- Cordis Corporation v. Boston Scientific Corporation et al.* (CA. No. 03-027-SLR), and *Boston Scientific Scimed, Inc. et al. v. Cordis Corporation et al.* (CA. No. 03-283-SLR), Deposition Transcript of Julio C. Palmaz.
- Arterial Vascular Engineering, Inc. v. Cordis Corporation, Johnson & Johnson and Expandable Grafts Partnership*, Cordis Corporation and Johnson & Johnson's Answer and Counterclaim, Feb. 27, 1998 (Civil Action No. 97-700-SLR).
- Plea Notes in *EPG et al. v. Boston Scientific et al.* in Netherlands (Sep. 12, 1997).
- Provisional Judgment *EPG et al. v. Boston Scientific et al.* in Netherlands (Oct. 29, 1997).
- Trial testimony transcripts from the Cordis Corporation et al. v. Medtronic AVE Inc., et al. liability trial dated Nov. 6-9, 13-17 and 20-21, 2000.
- Jury verdict form from the Cordis Corporation et al. v. Medtronic AVE, Inc. et al. liability trial.
- Hearing testimony transcript from the consolidated Cordis Corporation et al. v. Medtronic AVE, Inc. et al. and Boston Scientific Corporation et al. inequitable conduct hearing dated Feb. 7-9 and 12, 2001.
- Cordis Corporation v. Medtronic Ave., Inc. et al.* OPINION, 97-550-SLR, dated Mar. 28, 2002.
- Cordis Corporation v. Advanced Cardiovascular Systems, Inc. et al.* (CA. No. 97-550-SLR), *Medtronic Ave, Inc. v. Cordis Corporation et al.* (CA. No. 97-700-SLR), *Boston Scientific Corporation v. Athicon, Inc. et al.* (CA. No. 98-19-SLR), Expert Report of John T. Goolkasian, Esq.
- Cordis Corporation v. Advanced Cardiovascular Systems, Inc. et al.* (CA. No. 97-550-SLR), *Medtronic AVE, Inc. v. Cordis Corporation et al.* (CA. No. 97-700-SLR), *Boston Scientific Corporation v. Athicon, Inc. et al.* (CA. 98-19-SLR), Expert Report of John F. Witherspoon.
- "Microbial Conversion of Rapamycin," Kuhnt et al., *Enzyme and Microbial Technology*, vol. 21, pp. 405-412, 1997.
- "Inhibitory Effects of Rapamycin on Intimal Hyperplasia After PTCA," Badimon et al., *JACC*, Mar. 1998.

* cited by examiner

U.S. Patent

May 29, 2007

Sheet 1 of 2

US 7,223,286 B2

FIG. 1




FIG. 1a




FIG. 2a





FIG. 2b



U.S. Patent

May 29, 2007

Sheet 2 of 2

US 7,223,286 B2

FIG. 3a




FIG. 3b





FIG. 4



US 7,223,286 B2

1

**LOCAL DELIVERY OF RAPAMYCIN FOR
TREATMENT OF PROLIFERATIVE
SEQUELAE ASSOCIATED WITH PTCA
PROCEDURES, INCLUDING DELIVERY
USING A MODIFIED STENT**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

This application is a continuation of Ser. No. 10/408,328, filed Apr. 7, 2003, now issued as U.S. Pat. No. 6,808,536, which in turn is a continuation of application Ser. No. 09/874,117, filed Jun. 4, 2001, now issued as U.S. Pat. No. 6,585,764, which is a continuation of application Ser. No. 09/061,568, filed Apr. 16, 1998, now issued as U.S. Pat. No. 6,273,913, which in turn claims benefit of provisional application Ser. No. 60/044,692, filed Apr. 18, 1997.

FIELD OF THE INVENTION

Delivery of rapamycin locally, particularly from an intravascular stent, directly from micropores in the stent body or mixed or bound to a polymer coating applied on stent, to inhibit neointimal tissue proliferation and thereby prevent restenosis. This invention also facilitates the performance of the stent in inhibiting restenosis.

BACKGROUND OF THE INVENTION

Re-narrowing (restenosis) of an arteriosclerotic coronary artery after percutaneous transluminal coronary angioplasty (PTCA) occurs in 10–50% of patients undergoing this procedure and subsequently requires either further angioplasty or coronary artery bypass graft. While the exact hormonal and cellular processes promoting restenosis are still being determined, our present understanding is that the process of PTCA, besides opening the arteriosclerotically obstructed artery, also injures resident coronary arterial smooth muscle cells (SMC). In response to this injury, adhering platelets, infiltrating macrophages, leukocytes, or the smooth muscle cells (SMC) themselves release cell derived growth factors with subsequent proliferation and migration of medial SMC through the internal elastic lamina to the area of the vessel intima. Further proliferation and hyperplasia of intimal SMC and, most significantly, production of large amounts of extracellular matrix over a period of 3–6 months results in the filling in and narrowing of the vascular space sufficient to significantly obstruct coronary blood flow.

Several recent experimental approaches to preventing SMC proliferation have shown promise although the mechanisms for most agents employed are still unclear. Heparin is the best known and characterized agent causing inhibition of SMC proliferation both in vitro and in animal models of balloon angioplasty-mediated injury. The mechanism of SMC inhibition with heparin is still not known but may be due to any or all of the following: 1) reduced expression of the growth regulatory protooncogenes c-fos and c-myc, 2) reduced cellular production of tissue plasminogen activator; are 3) binding and dequstration of growth regulatory factors such as fibrovalent growth factor (FGF).

Other agents which have demonstrated the ability to reduce myointimal thickening in animal models of balloon vascular injury are angiopeptin (a somatostatin analog), calcium channel blockers, angiotensin converting enzyme inhibitors (captopril, cilazapril), cyclosporin A, trapidil (an antianginal, antiplatelet agent), terbinafine (antifungal),

2

colchicine and taxol (antitubulin antiproliferatives), and c-myc and c-myb antisense oligonucleotides.

Additionally, a goat antibody to the SMC mitogen platelet derived growth factor (PDGF) has been shown to be effective in reducing myointimal thickening in a rat model of balloon angioplasty injury, thereby implicating PDGF directly in the etiology of restenosis. Thus, while no therapy has as yet proven successful clinically in preventing restenosis after angioplasty, the in vivo experimental success of several agents known to inhibit SMC growth suggests that these agents as a class have the capacity to prevent clinical restenosis and deserve careful evaluation in humans.

Coronary heart disease is the major cause of death in men over the age of 40 and in women over the age of fifty in the western world. Most coronary artery-related deaths are due to atherosclerosis. Atherosclerotic lesions which limit or obstruct coronary blood flow are the major cause of ischemic heart disease related mortality and result in 500,000–600,000 deaths in the United States annually. To arrest the disease process and prevent the more advanced disease states in which the cardiac muscle itself is compromised, direct intervention has been employed via percutaneous transluminal coronary angioplasty (PTCA) or coronary artery bypass graft (CABG).

PTCA is a procedure in which a small balloon-tipped catheter is passed down a narrowed coronary artery and then expanded to re-open the artery. It is currently performed in approximately 250,000–300,000 patients each year. The major advantage of this therapy is that patients in which the procedure is successful need not undergo the more invasive surgical procedure of coronary artery bypass graft. A major difficulty with PTCA is the problem of post-angioplasty closure of the vessel, both immediately after PTCA (acute reocclusion) and in the long term (restenosis).

The mechanism of acute reocclusion appears to involve several factors and may result from vascular recoil with resultant closure of the artery and/or deposition of blood platelets along the damaged length of the newly opened blood vessel followed by formation of a fibrin/red blood cell thrombus. Recently, intravascular stents have been examined as a means of preventing acute reclosure after PTCA.

Restenosis (chronic reclosure) after angioplasty is a more gradual process than acute reocclusion: 30% of patients with subtotal lesions and 50% of patients with chronic total lesions will go on to restenosis after angioplasty. While the exact mechanism for restenosis is still under active investigation, the general aspects of the restenosis process have been identified.

In the normal arterial wall, smooth muscle cells (SMC) proliferate at a low rate (<0.1%/day; ref). SMC in vessel wall exists in a 'contractile' phenotype characterized by 80–90% of the cell cytoplasmic volume occupied with the contractile apparatus. Endoplasmic reticulum, golgi bodies, and free ribosomes are few and located in the perinuclear region. Extracellular matrix surrounds SMC and is rich in heparin-like glycosylaminoglycans which are believed to be responsible for maintaining SMC in the contractile phenotypic state.

Upon pressure expansion of an intracoronary balloon catheter during angioplasty, smooth muscle cells within the arterial wall become injured. Cell derived growth factors such as platelet derived growth factor (PDGF), basic fibroblast growth factor (bFGF), epidermal growth factor (EGF), etc. released from platelets (i.e., PDGF) adhering to the damaged arterial luminal surface, invading macrophages and/or leukocytes, or directly from SMC (i.e., bFGF) provoke a proliferation and migratory response in medial SMC.

US 7,223,286 B2

3

These cells undergo a phenotypic change from the contractile phenotype to a 'synthetic' phenotype characterized by only few contractile filament bundles but extensive rough endoplasmic reticulum, golgi and free ribosomes. Proliferation/migration usually begins within 1-2 days post-injury and peaks at 2 days in the media, rapidly declining thereafter (Campbell et al., In: *Vascular Smooth Muscle Cells in Culture*, Campbell, J. H. and Campbell, G. R., Eds, CRC Press, Boca Ration, 1987, pp. 39-55); Clowes, A. W. and Schwartz, S. M., *Circ. Res.* 56:139-145, 1985).

Finally, daughter synthetic cells migrate to the intimal layer of arterial smooth muscle and continue to proliferate. Proliferation and migration continues until the damaged luminal endothelial layer regenerates at which time proliferation ceases within the intima, usually within 7-14 days postinjury. The remaining increase in intimal thickening which occurs over the next 3-6 months is due to an increase in extracellular matrix rather than cell number. Thus, SMC migration and proliferation is an acute response to vessel injury while intimal hyperplasia is a more chronic response. (Liu et al., *Circulation*, 79:1374-1387, 1989).

Patients with symptomatic reocclusion require either repeat PTCA or CABG. Because 30-50% of patients undergoing PTCA will experience restenosis, restenosis has clearly limited the success of PTCA as a therapeutic approach to coronary artery disease. Because SMC proliferation and migration are intimately involved with the pathophysiological response to arterial injury, prevention of SMC proliferation and migration represents a target for pharmacological intervention in the prevention of restenosis.

SUMMARY OF THE INVENTION

Novel Features and Applications to Stent Technology

Currently, attempts to improve the clinical performance of stents have involved some variation of either applying a coating to the metal, attaching a covering or membrane, or embedding material on the surface via ion bombardment. A stent designed to include reservoirs is a new approach which offers several important advantages over existing technologies.

Local Drug Delivery from a Stent to Inhibit Restenosis

In this application, it is desired to deliver a therapeutic agent to the site of arterial injury. The conventional approach has been to incorporate the therapeutic agent into a polymer material which is then coated on the stent. The ideal coating material must be able to adhere strongly to the metal stent both before and after expansion, be capable of retaining the drug at a sufficient load level to obtain the required dose, be able to release the drug in a controlled way over a period of several weeks, and be as thin as possible so as to minimize the increase in profile. In addition, the coating material should not contribute to any adverse response by the body (i.e., should be non-thrombogenic, non-inflammatory, etc.). To date, the ideal coating material has not been developed for this application.

An alternative would be to design the stent to contain reservoirs which could be loaded with the drug. A coating or membrane of biocompatible material could be applied over the reservoirs which would control the diffusion of the drug from the reservoirs to the artery wall.

One advantage of this system is that the properties of the coating can be optimized for achieving superior biocompatibility and adhesion properties, without the addition requirement of being able to load and release the drug. The size,

4

shape, position, and number of reservoirs can be used to control the amount of drug, and therefore the dose delivered.

DESCRIPTION OF THE DRAWINGS

The invention will be better understood in connection with the following figures in which FIGS. 1 and 1A are top views and section views of a stent containing reservoirs as described in the present invention;

FIGS. 2a and 2b are similar views of an alternate embodiment of the stent with open ends;

FIGS. 3a and 3b are further alternate figures of a device containing a grooved reservoir; and

FIG. 4 is a layout view of a device containing a reservoir as in FIG. 3.

DETAILED DESCRIPTION OF THE INVENTION

Pharmacological attempts to prevent restenosis by pharmacologic means have thus far been unsuccessful and all involve systemic administration of the trial agents. Neither aspirin-dipyridamole, ticlopidine, acute heparin administration, chronic warfarin (6 months) nor methylprednisolone have been effective in preventing restenosis although platelet inhibitors have been effective in preventing acute reocclusion after angioplasty. The calcium antagonists have also been unsuccessful in preventing restenosis, although they are still under study. Other agents currently under study include thromboxane inhibitors, prostacyclin mimetics, platelet membrane receptor blockers, thrombin inhibitors and angiotensin converting enzyme inhibitors. These agents must be given systemically, however, and attainment of a therapeutically effective dose may not be possible; antiproliferative (or anti-restenosis) concentrations may exceed the known toxic concentrations of these agents so that levels sufficient to produce smooth muscle inhibition may not be reached (Lang et al., 42 *Ann. Rev. Med.*, 127-132 (1991); Popma et al., 84 *Circulation*, 1426-1436 (1991)).

Additional clinical trials in which the effectiveness for preventing restenosis of dietary fish oil supplements, thromboxane receptor antagonists, cholesterol lowering agents, and serotonin antagonists has been examined have shown either conflicting or negative results so that no pharmacological agents are as yet clinically available to prevent post-angioplasty restenosis (Franklin, S. M. and Faxon, D. P., 4 *Coronary Artery Disease*, 232-242 (1993); Serruys, P. W. et al., 88 *Circulation*, (part 1) 1588-1601, (1993).

Conversely, stents have proven useful in preventing reducing the proliferation of restenosis. Stents, such as the stent 10 seen in layout in FIG. 4, balloon-expandable slotted metal tubes (usually but not limited to stainless steel), which when expanded within the lumen of an angioplastied coronary artery, provide structural support to the arterial wall. This support is helpful in maintaining an open path for blood flow. In two randomized clinical trials, stents were shown to increase angiographic success after PTCA, increase the stenosed blood vessel lumen and to reduce the lesion recurrence at 6 months (Serruys et al., 331 *New Eng Jour. Med.* 495, (1994); Fischman et al., 331 *New Eng Jour. Med.* 496-501 (1994). Additionally, in a preliminary trial, heparin coated stents appear to possess the same benefit of reduction in stenosis diameter at follow-up as was observed with non-heparin coated stents. Additionally, heparin coating appears to have the added benefit of producing a reduction in sub-acute thrombosis after stent implantation (Serruys et al., 93 *Circulation*, 412-422, (1996). Thus, 1) sustained

US 7,223,286 B2

5

mechanical expansion of a stenosed coronary artery has been shown to provide some measure of restenosis prevention, and 2) coating of stents with heparin has demonstrated both the feasibility and the clinical usefulness of delivering drugs to local, injured tissue off the surface of the stent.

Numerous agents are being actively studied as antiproliferative agents for use in restenosis and have shown some activity in experimental animal models. These include: heparin and heparin fragments (Clowes and Karnovsky, 265 *Nature*, 25–626, (1977); Guyton, J. R. et al. 46 *Circ. Res.*, 625–634, (1980); Clowes, A. W. and Clowes, M. M., 52 *Lab. Invest.*, 611–616, (1985); Clowes, A. W. and Clowes, M. M., 58 *Circ. Res.*, 839–845 (1986); Majesky et al., 61 *Circ Res.*, 296–300, (1987); Snow et al., 137 *Am. J. Pathol.*, 313–330 (1990); Okada, T. et al., 25 *Neurosurgery*, 92–898, (1989) 15 colchicine (Currier, J. W. et al., 80 *Circulation*, 11–66, (1989), taxol (ref), angiotensin converting enzyme (ACE) inhibitors (Powell, J. S. et al., 245 *Science*, 186–188 (1989), angiopeptin (Lundergan, C. F. et al., 17 *Am. J. Cardiol. (Suppl. B)*; 132B–136B (1991), Cyclosporin A (Jonasson, L. et al., 85 *Proc. Natl. Acad. Sci.*, 2303 (1988), goat-anti-rabbit PDGF antibody (Ferns, G. A. A., et al., 253 *Science*, 1129–1132 (1991), terbinafine (Nemecek, G. M. et al., 248 *J. Pharmacol. Exp. Thera.*, 1167–11747 (1989), trapidil (Liu, M. W. et al., 81 *Circulation*, 1089–1093 (1990), interferon-gamma (Hansson, G. K. and Holm, 84 *J. Circulation*, 1266–1272 (1991), steroids (Colburn, M. D. et al., 15 *J. Vasc. Surg.*, 510–518 (1992), see also Berk, B. C. et al., 17 *J. Am. Coll. Cardiol.*, 111B–117B (1991), ionizing radiation (ref), fusion toxins (ref) antisense oligonucleotides (ref), 20 gene vectors (ref), and rapamycin (see below).

Of particular interest in rapamycin. Rapamycin is a macrolide antibiotic which blocks IL-2-mediated T-cell proliferation and possesses antiinflammatory activity. While the precise mechanism of rapamycin is still under active investigation, rapamycin has been shown to prevent the G₁ to S phase progression of T-cells through the cell cycle by inhibiting specific cell cyclins and cyclin-dependent protein kinases (Siekierka, *Immunol. Res.* 13: 110–116, 1994). The antiproliferative action of rapamycin is not limited to T-cells; Marx et al. (*Circ Res* 76:412–417, 1995) have demonstrated that rapamycin prevents proliferation of both rat and human SMC in vitro while Poon et al. have shown the rat, porcine, and human SMC migratin can also be inhibited by rapamycin (*J Clin Invest* 98: 2277–2283, 1996). 25 Thus, rapamycin is capable of inhibiting both the inflammatory response known to occur after arterial injury and stent implantation, as well as the SMC hyperproliferative response. In fact, the combined effects of rapamycin have been demonstrated to result in a diminished SMC hyperproliferative response in a rat femoral artery graft model and in both rat and porcine arterial balloon injury models (Gregory et al., *Transplantation* 55:1409–1418, 1993; Gallo et al., in press, (1997)). These observations clearly support the potential use of rapamycin in the clinical setting of post-angioplasty restenosis.

Although the ideal agent for restenosis has not yet been identified, some desired properties are clear: inhibition of local thrombosis without the risk systemic bleeding complications and continuous and prevention of the dequale of arterial injury, including local inflammation and sustained prevention smooth muscle proliferation at the site of angioplasty without serious systemic complications. Inasmuch as stents prevent at least a portion of the restenosis process, an agent which prevents inflammation and the proliferation of SMC combined with a stent may provide the most efficacious treatment for post-angioplasty restenosis.

6

Experiments

Agents: Rapamycin (sirolimus) structural analogs (macrocyclic lactones) and inhibitors of cell-cycle progression.

Delivery Methods:

These can vary:

Local delivery of such agents (rapamycin) from the struts of a stent, from a stent graft, grafts, stent cover or sheath.

Involving comixture with polymers (both degradable and nondegrading) to hold the drug to the stent or graft.

or entrapping the drug into the metal of the stent or graft body which has been modified to contain micropores or channels, as will be explained further herein.

or including covalent binding of the drug to the stent via solution chemistry techniques (such as via the Carmeda process) or dry chemistry techniques (e.g. vapour deposition methods such as rf-plasma polymerization) and combinations thereof.

Catheter delivery intravascularly from a tandem balloon or a porous balloon for intramural uptake

Extravascular delivery by the pericardial route

Extravascular delivery by the adventitial application of sustained release formulations.

Uses: for inhibition of cell proliferation to prevent neointimal proliferation and restenosis.

prevention of tumor expansion from stents

prevent ingrowth of tissue into catheters and shunts inducing their failure.

1. Experimental Stent Delivery Method—Delivery from Polymer Matrix:

Solution of Rapamycin, prepared in a solvent miscible with polymer carrier solution, is mixed with solution of polymer at final concentration range 0.001 weight % to 30 weight % of drug. Polymers are biocompatible (i.e., not elicit any negative tissue reaction or promote mural thrombus formation) and degradable, such as lactone-based polyesters or copolyesters, e.g., polylactide, polycaprolactone-glycolide, polyorthoesters, polyanhydrides; poly-amino acids; polysaccharides; polyphosphazenes; poly(ether-ester) copolymers, e.g., PEO-PLLA, or blends thereof. Nonabsorbable biocompatible polymers are also suitable candidates. Polymers such as polydimethylsiloxane; poly(ethylene-vinylacetate); acrylate based polymers or copolymers, e.g., poly(hydroxyethyl methylmethacrylate, polyvinyl pyrrolidone; fluorinated polymers such as polytetrafluoroethylene; cellulose esters.

Polymer/drug mixture is applied to the surfaces of the stent by either dip-coating, or spray coating, or brush coating or dip/spin coating or combinations thereof, and the solvent allowed to evaporate to leave a film with entrapped rapamycin.

2. Experimental Stent Delivery Method—Delivery from Microporous Depots in Stent Through a Polymer Membrane Coating:

Stent, whose body has been modified to contain micropores or channels is dipped into a solution of Rapamycin, range 0.001 wt % to saturated, in organic solvent such as acetone or methylene chloride, for sufficient time to allow solution to permeate into the pores. (The dipping solution can also be compressed to improve the loading efficiency.) After solvent has been allowed to evaporate, the stent is dipped briefly in fresh solvent to remove excess surface bound drug. A solution of polymer, chosen from any identified in the first experimental method, is applied to the stent as detailed above. This outer layer of polymer will act as diffusion-controller for release of drug.

US 7,223,286 B2

7

3. Experimental Stent Delivery Method—Delivery via Lysis of a Covalent Drug Tether

Rapamycin is modified to contain a hydrolytically or enzymatically labile covalent bond for attaching to the surface of the stent which itself has been chemically derivatized to allow covalent immobilization. Covalent bonds such as ester, amides or anhydrides may be suitable for this.

4. Experimental Method—Pericardial Delivery

A: Polymeric Sheet Rapamycin is combined at concentration range previously highlighted, with a degradable polymer such as poly(caprolactone-glycolide) or non-degradable polymer, e.g., polydimethylsiloxane, and mixture cast as a thin sheet, thickness range 10 μ to 1000 μ . The resulting sheet can be wrapped perivascularly on the target vessel. Preference would be for the absorbable polymer.

B: Conformal Coating: Rapamycin is combined with a polymer that has a melting temperature just above 37° C., range 40°–45° C. Mixture is applied in a molten state to the external side of the target vessel. Upon cooling to body temperature the mixture solidifies conformably to the vessel wall. Both non-degradable and absorbable biocompatible polymers are suitable.

As seen in the figures it is also possible to modify currently manufactured stents in order to adequately provide the drug dosages such as rapamycin. As seen in FIGS. 1a, 2a and 3a, any stent strut 10, 20, 30 can be modified to have a certain reservoir or channel 11, 21, 31. Each of these reservoirs can be open or closed as desired. These reservoirs can hold the drug to be delivered. FIG. 4 shows a stent 40 with a reservoir 45 created at the apex of a flexible strut. Of course, this reservoir 45 is intended to be useful to deliver rapamycin or any other drug at a specific point of flexibility of the stent. Accordingly, this concept can be useful for "second generation" type stents.

In any of the foregoing devices, however, it is useful to have the drug dosage applied with enough specificity and enough concentration to provide an effective dosage in the lesion area. In this regard, the reservoir size in the stent struts must be kept at a size of about 0.0005" to about 0.003". Then, it should be possible to adequately apply the drug dosage at the desired location and in the desired amount.

These and other concepts will be disclosed herein. It would be apparent to the reader that modifications are possible to the stent or the drug dosage applied. In any event, however, the any obvious modifications should be perceived to fall within the scope of the invention which is to be realized from the attached claims and their equivalents.

What is claimed is:

1. A stent having a coating applied thereto, wherein said coating comprises a biocompatible polymer/drug mixture and said drug is rapamycin or a macrocyclic lactone analog thereof.

2. A stent according to claim 1 comprising a generally thin walled cylinder containing a plurality of generally solid struts to which said coating is applied.

3. A stent according to claim 2 further comprising a channel formed in at least one of said struts.

4. A stent according to claim 3, wherein said channel has a closed perimeter on all sides, an open top and a generally rectangular perimeter, and said channel is smaller in all dimensions than said strut.

5. A stent according to claim 1 wherein the coating is dip-coated onto the stent.

6. A stent according to claim 1 wherein the coating is spray-coated onto the stent.

8

7. A stent according to claim 1 wherein said rapamycin or macrocyclic lactone analog thereof is contained in the coating at a weight percentage of about 30%.

8. A stent according to claim 1 wherein the coating comprises a degradable polymer.

9. A stent according to claim 1 wherein the coating comprises a nonabsorbable polymer.

10. A stent according to claim 1 wherein the coating comprises a lactone-based polyester; a lactone-based copolyester; a polyanhydride; a polyaminoacid; a polysaccharide; a polyphosphazene; a poly(ether-ester) copolymer; a polydimethylsiloxane; a poly(ethylene)vinylacetate; a poly(hydroxy)ethylmethacrylate; an acrylate based polymer; an acrylate based copolymer; a polyvinyl pyrrolidone; a cellulose ester; a fluorinated polymer; or a blend thereof.

11. A stent according to claim 10 wherein the coating comprises a lactone-based polyester.

12. A stent according to claim 10 wherein the coating comprises a lactone-based copolyester.

13. A stent according to claim 10 wherein the coating comprises a polyanhydride.

14. A stent according to claim 10 wherein the coating comprises a polyaminoacid.

15. A stent according to claim 10 wherein the coating comprises a polysaccharide.

16. A stent according to claim 10 wherein the coating comprises a polyphosphazene.

17. A stent according to claim 10 wherein the coating comprises a poly(ether-ester) copolymer.

18. A stent according to claim 10 wherein the coating comprises a polydimethylsiloxane.

19. A stent according to claim 10 wherein the coating comprises a poly(ethylene)vinylacetate.

20. A stent according to claim 10 wherein the coating comprises a poly(hydroxy)ethylmethacrylate.

21. A stent according to claim 10 wherein the coating comprises an acrylate based polymer.

22. A stent according to claim 10 wherein the coating comprises an acrylate based copolymer.

23. A stent according to claim 10 wherein the coating comprises a polyvinyl pyrrolidone.

24. A stent according to claim 10 wherein the coating comprises a cellulose ester.

25. A stent according to claim 10 wherein the coating comprises a fluorinated polymer.

26. A stent according to claim 10 wherein the fluorinated polymer is polytetrafluoroethylene.

27. A stent according to any one of claims 1 to 26 wherein said drug is a macrocyclic lactone analog of rapamycin.

28. A stent according to any one of claims 1 to 26 that provides a controlled release of said rapamycin or macrocyclic lactone analog thereof over a period of several weeks.

29. A stent according to claim 28 wherein said drug is a macrocyclic lactone analog of rapamycin.

30. A stent according to any one of claims 1 to 26 that releases said rapamycin or macrocyclic lactone analog thereof over a period of at least 14 days.

31. A stent according to claim 30 wherein said drug is a macrocyclic lactone analog of rapamycin.

32. A stent according to any one of claims 1 to 26 wherein said rapamycin or macrocyclic lactone analog thereof is present in a therapeutically beneficial amount to inhibit neointimal proliferation.

33. A stent according to claim 32 wherein said drug is a macrocyclic lactone analog of rapamycin.

US 7,223,286 B2

9

34. A stent according to claim 33 that releases said macrocyclic lactone analog of rapamycin over a period of at least 14 days.

35. A stent according to claim 34 wherein the coating comprises a fluorinated polymer.

36. A stent according to claim 35 wherein the coating further comprises an acrylate based polymer or copolymer.

37. A stent according to claim 33 that provides a controlled release of said rapamycin or macrocyclic lactone analog thereof over a period of several weeks.

38. A stent according to claim 37 wherein the coating comprises a fluorinated polymer.

39. A stent according to claim 38 wherein the coating further comprises an acrylate based polymer or copolymer.

40. A device comprising a metallic stent, a biocompatible polymeric carrier and a drug, wherein said drug is rapamycin or a macrocyclic lactone analog thereof and is present in an amount effective to inhibit neointimal proliferation.

41. A device according to claim 40 wherein said polymeric carrier and drug are mixed together.

42. A device according to claim 40 wherein said polymeric carrier is bound to the drug.

43. A device according to claim 40 wherein the drug is entrapped on the surface of the stent by said polymeric carrier.

44. A device according to claim 40 wherein said stent comprises a generally thin walled cylinder containing a plurality of generally solid struts to which said polymeric carrier and drug are applied.

45. A device according to claim 44 further comprising a channel formed in at least one of said struts.

46. A device according to claim 45, wherein said channel has a closed perimeter on all sides, an open top and a generally rectangular perimeter, and said channel is smaller in all dimensions than said strut.

47. A device according to claim 40 wherein the polymeric carrier and drug are dip-coated onto the stent.

48. A device according to claim 40 wherein the polymeric carrier and drug are spray-coated onto the stent.

49. A device according to claim 40 wherein the weight ratio of drug to polymeric carrier is about 3:7.

50. A device according to claim 40 wherein the polymeric carrier comprises a degradable polymer.

51. A device according to claim 40 wherein the polymeric carrier comprises a nonabsorbable polymer.

52. A device according to claim 40 wherein the polymeric carrier comprises a lactone-based polyester; a lactone-based copolyester; a polyanhydride; a polyaminoacid; a polysaccharide; a polyphosphazene; a poly(ether-ester) copolymer; a polydimethylsiloxane; a poly(ethylene)vinylacetate; a poly(hydroxy)ethylmethacrylate; an acrylate based polymer; an acrylate based copolymer; a polyvinyl pyrrolidone; a cellulose ester; a fluorinated polymer; or a blend thereof.

53. A device according to claim 52 wherein the polymeric carrier comprises a lactone-based polyester.

10

54. A device according to claim 52 wherein the polymeric carrier comprises a lactone-based copolyester.

55. A device according to claim 52 wherein the polymeric carrier comprises a polyanhydride.

56. A device according to claim 52 wherein the polymeric carrier comprises a polyaminoacid.

57. A device according to claim 52 wherein the polymeric carrier comprises a polysaccharide.

58. A device according to claim 52 wherein the polymeric carrier comprises a polyphosphazene.

59. A device according to claim 52 wherein the polymeric carrier comprises a poly(ether-ester) copolymer.

60. A device according to claim 52 wherein the polymeric carrier comprises a polydimethylsiloxane.

61. A device according to claim 52 wherein the polymeric carrier comprises a poly(ethylene)vinylacetate.

62. A device according to claim 52 wherein the polymeric carrier comprises a poly(hydroxy)ethylmethacrylate.

63. A device according to claim 52 wherein the polymeric carrier comprises an acrylate based polymer.

64. A device according to claim 52 wherein the polymeric carrier comprises an acrylate based copolymer.

65. A device according to claim 52 wherein the polymeric carrier comprises a polyvinyl pyrrolidone.

66. A device according to claim 52 wherein the polymeric carrier comprises a cellulose ester.

67. A device according to claim 52 wherein the polymeric carrier comprises a fluorinated polymer.

68. A device according to claim 67 wherein the fluorinated polymer is polytetrafluoroethylene.

69. A device according to any one of claims 40 to 68 wherein said drug is a macrocyclic lactone analog of rapamycin.

70. A device according to any one of claims 40 to 68 that provides a controlled release of said rapamycin or macrocyclic lactone analog thereof over a period of several weeks.

71. A device according to claim 70 wherein said drug is a macrocyclic lactone analog of rapamycin.

72. A device according to claim 71 wherein the polymeric carrier comprises a fluorinated polymer.

73. A device according to claim 72 wherein the polymeric carrier further comprises an acrylate based polymer or copolymer.

74. A device according to any one of claims 40 to 68 that releases said drug over a period of at least 14 days.

75. A device according to claim 74 wherein said drug is a macrocyclic lactone analog of rapamycin.

76. A device according to claim 75 wherein the polymeric carrier comprises a fluorinated polymer.

77. A device according to claim 76 wherein the polymeric carrier further comprises an acrylate based polymer or copolymer.

* * * * *